arXiv:1203.1843v1 [math.CV] 8 Mar 2012

QUANTITATIVE EQUIDISTRIBUTION FOR THE SOLUTIONS OF A SYSTEM OF SPARSE POLYNOMIAL EQUATIONS

CARLOS D'ANDREA, ANDRÉ GALLIGO, AND MARTÍN SOMBRA

Abstract. For a system of Laurent polynomials $f_1, \dots, f_n \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ whose coefficients are not too big with respect to its facet resultants, we show that the solutions in the algebraic torus $(\mathbb{C}^\times)^n$ of the system of equations $f_1 = \dots = f_n = 0$, are approximately equidistributed near the unit polycircle. This generalizes to the multivariate case, up to a exponent, a classical result due to Erdös and Turán on the distribution of the arguments of the roots of a univariate polynomial.

We apply this result to bound the number of real roots of a system of Laurent polynomials, and to study the asymptotic distribution of the roots of systems of Laurent polynomials over $\mathbb{Z}$ and of random systems of Laurent polynomials over $\mathbb{C}$.

1. Introduction and statement of results

A celebrated result due to Erdös and Turán says that, for a univariate polynomial over $\mathbb{C}$ whose middle coefficients are not too big with respect to the extremal ones, the arguments of its roots are approximately equidistributed [ET50]. Combined with a recent result of Hughes and Nikeghbali [HN08], this shows that the roots of such a polynomial cluster near the unit circle.

We introduce some notation to precise this result. Let Z be an effective cycle of $\mathbb{C}^\times$ of dimension 0, that is, a formal finite sum

$$Z = \sum_{\xi} m_\xi \xi$$

with $\xi \in \mathbb{C}^\times$ and $m_\xi \in \mathbb{N}$. The degree of Z, denoted $\deg(Z)$, is defined as the sum of its multiplicities m_ξ. We assume that $Z \neq 0$ or, equivalently, that $\deg(Z) \geq 1$.

For each $-\pi \leq \alpha < \beta \leq \pi$, consider the cycle

$$Z_{\alpha,\beta} = \sum_{\alpha < \arg(\xi) \leq \beta} m_\xi \xi,$$

where $\arg(\xi)$ denotes the argument of ξ. The *angle discrepancy* of Z is defined as

$$\Delta_{\mathrm{ang}}(Z) = \sup_{-\pi \leq \alpha < \beta \leq \pi} \left| \frac{\deg(Z_{\alpha,\beta})}{\deg(Z)} - \frac{\beta - \alpha}{2\pi} \right|.$$

For $0 < \varepsilon < 1$, consider also the cycle

$$Z_\varepsilon = \sum_{1-\varepsilon < |\xi| < (1-\varepsilon)^{-1}} m_\xi \xi.$$

Date: February 25, 2013.

2010 *Mathematics Subject Classification.* Primary 30C15; Secondary 11K38, 13P15, 42B05.

Key words and phrases. Equidistribution, zeros of systems of polynomials, exponential sums, sparse resultants, random systems of polynomials.

D'Andrea was partially supported by the MICINN research project MTM2010-20279 (Spain). Galligo was partially supported by the European Marie Curie network SAGA. Sombra was partially supported by the MICINN research project MTM2009-14163-C02-01 (Spain).

The *radius discrepancy* of Z with respect to ε is defined as

$$\Delta_{\mathrm{rad}}(Z,\varepsilon) = 1 - \frac{\deg(Z_\varepsilon)}{\deg(Z)}.$$

For a polynomial $f \in \mathbb{C}[x] \setminus \{0\}$, we denote by $Z(f)$ the 0-dimensional effective cycle of $\mathbb{C}^\times$ defined by its roots and their corresponding multiplicities. We also set $\|f\|_{\sup} = \sup_{|z|=1} |f(z)|$.

Theorem 1.1. *Let* $f = a_0 + \cdots + a_d x^d \in \mathbb{C}[x]$ *with* $d \geq 1$ *and* $a_0 a_d \neq 0$, *and* $0 < \varepsilon < 1$. *Then*

$$\Delta_{\mathrm{ang}}(Z(f)) \leq c\sqrt{\frac{1}{d}\log\left(\frac{\|f\|_{\sup}}{\sqrt{|a_0 a_d|}}\right)}, \quad \Delta_{\mathrm{rad}}(Z(f),\varepsilon) \leq \frac{2}{\varepsilon d}\log\left(\frac{\|f\|_{\sup}}{\sqrt{|a_0 a_d|}}\right),$$

with $c = \sqrt{2\pi/\kappa} = 2.5619\ldots$, *where* $\kappa = \sum_{m=0}^{\infty} \frac{(-1)^{m-1}}{(2m+1)^2} = 0.915965594\ldots$ *is Catalan's constant.*

The more interesting (and hardest) part is the bound for the angle discrepancy. The original Erdös-Turán result states [ET50]

$$\Delta_{\mathrm{ang}}(Z(f)) \leq 16\sqrt{\frac{1}{d}\log\left(\frac{\sum_j |a_j|}{\sqrt{|a_0 a_d|}}\right)}.$$

A few years after that paper, Ganelius [Gan54] replaced the ℓ^1-norm $\sum_j |a_j|$ by the smaller quantity $\|f\|_{\sup}$ and improved the value of the constant to $c \leq \sqrt{2\pi/\kappa}$. On the other hand, Amoroso and Mignotte [AM96] showed that the optimal value of c cannot be smaller than $\sqrt{2}$. The bound for the radius discrepancy is due to Hughes and Nikeghbali [HN08].

In this text, we study the distribution of the solutions in the algebraic torus $(\mathbb{C}^\times)^n$ of a system of multivariate polynomial equations. For instance, consider the following system of bivariate polynomials:

$$f_1 = x_1^{13} + x_1 x_2^{12} + x_2^{13} + 1, \quad f_2 = x_1^{12} x_2 - x_2^{13} - x_1 x_2 + 1 \quad \in \mathbb{C}[x_1, x_2]. \tag{1.2}$$

These are polynomials with a moderate degree and small integer coefficients. By direct computation, we can verify that the solutions in $(\mathbb{C}^\times)^2$ of the system of equations $f_1 = f_2 = 0$ are aproximately equidistributed near the unit polycircle $S^1 \times S^1$ (Figure 1). This example and others of the same kind suggest that Theorem 1.1 has an extension to higher dimensions.



FIGURE 1. Angle and radius distribution of the zeros of the system (1.2)

The study of the distribution of the solutions of a system of multivariate polynomial equations has been addressed from different perspectives. In the frame of his theory of fewnomials, Khovanskii has shown that one can estimate the distribution of the arguments of these solutions in terms of the number of monomials and the Newton polytopes of the input system, see for instance [Kho91, §3.13, Theorem 2]. There are also several interesting results by Shiffman, Zelditch and Bloom on the asymptotic distribution of the solutions of a random system of polynomial equations, see for instance [SZ04, BS07] and the references therein.

Our purpose in this text is to obtain an extension of Theorem 1.1 to systems of Laurent polynomials with a given support. For $i = 1, \dots, n$, let $\mathcal{A}_i$ be a non-empty finite subset of $\mathbb{Z}^n$ and $Q_i = \operatorname{conv}(\mathcal{A}_i) \subset \mathbb{R}^n$ its convex hull. Set $D = \operatorname{MV}(Q_1, \dots, Q_n)$ for the mixed volume of these lattice polytopes, and assume that $D \geq 1$. For each i, let f_i be a Laurent polynomial with support contained in $\mathcal{A}_i$, that is,

$$f_i = \sum_{\boldsymbol{a} \in \mathcal{A}_i} \alpha_{i,\boldsymbol{a}}\, \boldsymbol{x}^{\boldsymbol{a}} \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$$

with $\alpha_{i,\boldsymbol{a}} \in \mathbb{C}$ and $\boldsymbol{x}^{\boldsymbol{a}} = x_1^{a_1} \dots x_n^{a_n}$ for each $\boldsymbol{a} = (a_1, \dots, a_n) \in \mathcal{A}_i$. Write $\boldsymbol{f} = (f_1, \dots, f_n)$ for short. For each primitive vector $\boldsymbol{v} \in \mathbb{Z}^n$, consider the corresponding "facet resultant", denoted by $\operatorname{Res}_{\mathcal{A}_1^{\boldsymbol{v}}, \dots, \mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}}) \in \mathbb{C}$, of $\boldsymbol{f}$ in the direction of $\boldsymbol{v}$ (Definition 3.4). These resultants can differ from 1 only for a finite number of $\boldsymbol{v}$'s. We assume that all of them are nonzero. This condition holds for a generic choice of $\boldsymbol{f}$ in the space of coefficients and, by Bernstein-Kushnirenko-Khovanskii's theorem [Ber75, Theorem B], it implies that all the solutions of $f_1 = \dots = f_n = 0$ are isolated and that their number, counted with multiplicities, is equal to D.

For a vector $\boldsymbol{w} \in S^{n-1}$ in the unit sphere of $\mathbb{R}^n$, we denote by $\boldsymbol{w}^{\perp} \subset \mathbb{R}^n$ its orthogonal subspace and by $\pi_{\boldsymbol{w}} \colon \mathbb{R}^n \to \boldsymbol{w}^{\perp}$ the orthogonal projection onto it. We denote by $\operatorname{MV}_{\boldsymbol{w}^{\perp}}$ the mixed volume of convex bodies of $\boldsymbol{w}^{\perp}$ induced by the Euclidean measure on $\boldsymbol{w}^{\perp}$ and, for $i = 1, \dots, n$, we set

$$D_{\boldsymbol{w},i} = \operatorname{MV}_{\boldsymbol{w}^{\perp}}\left(\pi_{\boldsymbol{w}}(Q_1), \dots, \pi_{\boldsymbol{w}}(Q_{i-1}), \pi_{\boldsymbol{w}}(Q_{i+1}), \dots, \pi_{\boldsymbol{w}}(Q_n)\right).$$

We then define the *Erdös-Turán size* of $\boldsymbol{f}$ as

$$\eta(\boldsymbol{f}) = \frac{1}{D} \sup_{\boldsymbol{w} \in S^{n-1}} \log \left(\frac{\prod_{i=1}^{n} \|f_i\|_{\sup}^{D_{\boldsymbol{w},i}}}{\prod_{\boldsymbol{v}} |\operatorname{Res}_{\mathcal{A}_1^{\boldsymbol{v}}, \dots, \mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}})|^{\frac{|\langle \boldsymbol{v}, \boldsymbol{w} \rangle|}{2}}} \right), \tag{1.3}$$

where the second product is taken over all primitive vectors $\boldsymbol{v} \in \mathbb{Z}^n$ and $\langle \cdot, \cdot \rangle$ denotes the standard inner product of $\mathbb{R}^n$. This is a generalization to the multivariate case of the quantity $\frac{1}{d} \log \left(\frac{\|f\|_{\sup}}{\sqrt{|a_0 a_d|}} \right)$ that appears in Theorem 1.1 (Proposition 3.12).

We set $Z(\boldsymbol{f})$ for the 0-dimensional effective cycle of $(\mathbb{C}^{\times})^n$ defined by the roots of $\boldsymbol{f}$ and their multiplicities. The angle and radius discrepancies of cycles of $(\mathbb{C}^{\times})^n$ are the obvious generalization of those for the univariate case (see Definition 2.1).

The following is our main result in this text.

Theorem 1.4. *Let $\mathcal{A}_1, \dots, \mathcal{A}_n$ be family of non-empty finite subsets of $\mathbb{Z}^n$ such that $\operatorname{MV}(Q_1, \dots, Q_n) \geq 1$ with $Q_i = \operatorname{conv}(\mathcal{A}_i)$. Let $f_1, \dots, f_n \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ with $\operatorname{supp}(f_i) \subset \mathcal{A}_i$ and such that $\operatorname{Res}_{\mathcal{A}_1^{\boldsymbol{v}}, \dots, \mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}}) \neq 0$ for all $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$. Then*

$$\Delta_{\mathrm{ang}}(Z(\boldsymbol{f})) \leq 66\, n\, 2^n (18 + \log^{+}(\eta(\boldsymbol{f})^{-1}))^{\frac{2}{3}(n-1)} \eta(\boldsymbol{f})^{\frac{1}{3}}.$$

with $\log^+(x) = \log(\max(1,x))$ *for* $x > 0$*. Also, for* $0 < \varepsilon < 1$,

$$\Delta_{\mathrm{rad}}(Z(\boldsymbol{f}), \varepsilon) \le \frac{2n}{\varepsilon}\eta(\boldsymbol{f}).$$

For $n = 1$, our bound for the angle discrepancy is worse than Erdös-Turán's, due to our approach. By analogy with the one-dimensional case, it is natural to ask if, in the setting of our result, a stronger inequality of the form

$$\Delta_{\mathrm{ang}}(Z(\boldsymbol{f})) \le c(n)\, \eta(\boldsymbol{f})^{\frac{1}{2}}$$

holds, with $c(n) > 0$ not depending on $\boldsymbol{f}$. It would be interesting to settle this question.

Theorem 1.4 has several consequences. For instance, we can derive from it a bound for the number of positive real solutions of a system of polynomial equations, in terms of its Erdös-Turán size. For a cycle $Z = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \boldsymbol{\xi}$ of $(\mathbb{C}^\times)^n$, set

$$Z_+ = \sum_{\boldsymbol{\xi} \in (\mathbb{R}_{>0})^n} m_{\boldsymbol{\xi}} \boldsymbol{\xi}.$$

Corollary 1.5. *Let notations be as in Theorem 1.4. Then*

$$\deg(Z(\boldsymbol{f})_+) \le 66\, n\, 2^n (18 + \log^+(\eta(\boldsymbol{f})^{-1}))^{\frac{2}{3}(n-1)} \eta(\boldsymbol{f})^{\frac{1}{3}} \deg(Z(\boldsymbol{f})).$$

We can also apply our result to study the asymptotic distribution of the roots of a sequence of systems of polynomials over $\mathbb{Z}$ with growing supports and whose coefficients are not too big. To be more precise, let Q_i, $i = 1, \dots, n$, be lattice polytopes in $\mathbb{R}^n$ such that $\mathrm{MV}(Q_1, \dots, Q_n) \ge 1$. For each $d \ge 1$ and $i = 1, \dots, n$, consider the finite subset of $\mathbb{Z}^n$ given by

$$\mathcal{A}_{i,d} = dQ_i \cap \mathbb{Z}^n. \tag{1.6}$$

For a Laurent polynomial $f \in \mathbb{C}[x_1^{\pm1}, \dots, x_n^{\pm1}]$, we denote by $\mathrm{supp}(f)$ its support, defined as the subset of $\mathbb{Z}^n$ of the exponents in its monomial terms. We also set $\|f\|_{\sup} = \sup_{|w_1|=1,\dots,|w_n|=1} |f(w_1, \dots, w_n)|$.

For a nonzero cycle $Z = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \boldsymbol{\xi}$ of $(\mathbb{C}^\times)^n$, we consider the discrete probability measure on $(\mathbb{C}^\times)^n$ defined by

$$\delta_Z = \frac{1}{\deg(Z)} \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \delta_{\boldsymbol{\xi}},$$

where $\delta_{\boldsymbol{\xi}}$ is the Dirac measure supported on the point $\boldsymbol{\xi}$. Let ν be the measure on $(\mathbb{C}^\times)^n$ supported on $(S^1)^n$ and whose restriction to this polycircle coincides with its Haar measure of total mass 1.

Recall that a sequence of measures $(\nu_d)_{d\ge1}$ on $(\mathbb{C}^\times)^n$ converges weakly to ν if, for every continuous function with compact support $h\colon (\mathbb{C}^\times)^n \to \mathbb{R}$, it holds

$$\lim_{d\to\infty} \int_{(\mathbb{C}^\times)^n} h \,\mathrm{d}\nu_d = \int_{(\mathbb{C}^\times)^n} h \,\mathrm{d}\nu.$$

If this is the case, we write $\lim_{d\to\infty} \nu_d = \nu$.

Theorem 1.7. *For* $d \ge 1$*, let* $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ *be a family of Laurent polynomials in* $\mathbb{Z}[x_1^{\pm1}, \dots, x_n^{\pm1}]$ *such that* $\mathrm{supp}(f_{i,d}) \subset dQ_i$, $\log\|f_{i,d}\|_{\sup} = o(d)$*, and* $\mathrm{Res}_{\mathcal{A}_{1,d}^{\boldsymbol{v}},\dots,\mathcal{A}_{n,d}^{\boldsymbol{v}}}(f_{1,d}^{\boldsymbol{v}}, \dots, f_{n,d}^{\boldsymbol{v}}) \ne 0$ *for all* $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$*. Then*

$$\lim_{d\to\infty} \delta_{Z(\boldsymbol{f}_d)} = \nu.$$

This result admits a quantitive version, which we state in Proposition 4.2. Theorem 1.7 is related to Bilu's equidistribution theorem for the Galois orbit of algebraic points in $(\mathbb{C}^\times)^n$ of small height [Bil97] which, at least for $n = 1$, also admits quantitative versions [Pet05, FRL06]. Indeed, when the cycles $Z(\boldsymbol{f}_d)$ are irreducible over $\mathbb{Q}$ or when, at least, they have a very large irreducible component, it is possible to deduce Theorem 1.7 from Bilu's theorem and the arithmetic Bézout theorem.

We can also apply Theorem 1.4 to study the distribution of the roots of a random system of Laurent polynomials over $\mathbb{C}$. We will show that, under some general conditions and without assuming any independence or equidistribution condition on the coefficients of the system, these roots tend to cluster uniformly near the unit polycircle.

Let us keep notations as above and set $\boldsymbol{\mathcal{A}}_d = (\mathcal{A}_{1,d}, \dots, \mathcal{A}_{n,d})$ with $\mathcal{A}_{i,d}$ as in (1.6). Each point of the projective space $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ can be identified with a system $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ of Laurent polynomials such that $\mathrm{supp}(f_{i,d}) \subset dQ_i$, $i = 1, \dots, n$, modulo a multiplicative scalar. The associated cycle $Z(\boldsymbol{f}_d)$ is well-defined, since it does not depend on this multiplicative scalar.

Let μ_d be the normalized Fubini-Study measure on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ of total mass 1, and g_d a probability density function on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ (see §4 for details). Let $\boldsymbol{f}_d$ be a random system of Laurent polynomials with $\mathrm{supp}(f_{i,d}) \subset dQ_i$, $i = 1, \dots, n$, distributed according to the probability law given by g_d with respect to μ_d. The *expected zero density measure* of $\boldsymbol{f}_d$ is the measure on $(\mathbb{C}^\times)^n$ defined, for a Borel subset U, as

$$\mathbb{E}(Z(\boldsymbol{f}_d); g_d)(U) = \int_{\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})} \deg(Z(\boldsymbol{f}_d)|_U)\, g_d(\boldsymbol{f}_d)\, \mathrm{d}\mu_d,$$

where $Z(\boldsymbol{f}_d)|_U$ denotes the cycle $\sum_{\boldsymbol{\xi} \in V(\boldsymbol{f}_d)_0 \cap U} m_{\boldsymbol{\xi}} \boldsymbol{\xi}$.

Theorem 1.8. *For $d \geq 1$, let g_d be a probability density function on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ with respect to the measure μ_d, and $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ a random system of Laurent polynomials with* $\mathrm{supp}(f_{i,d}) \subset dQ_i$, *$i = 1, \dots, n$, distributed according to the probability law given by g_d. Assume that the sequence $(g_d)_{d \geq 1}$ is uniformly bounded. Then*

$$\lim_{d \to \infty} \frac{\mathbb{E}(Z(\boldsymbol{f}_d); g_d)}{d^n\, \mathrm{MV}(Q_1, \dots, Q_n)} = \nu.$$

As an application, consider a random system of Laurent polynomials $\boldsymbol{f}_d$ with $\mathrm{supp}(f_{i,d}) \subset dQ_i$ whose coefficients are independent complex Gaussian random variables with mean 0 and variance 1. The random cycle $Z(\boldsymbol{f}_d)$ might be described by the uniform distribution on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}})$ (see Example 4.14 for details). Then, Theorem 1.8 implies that the roots of $\boldsymbol{f}_d$ converge weakly to the equidistribution on $(S^1)^n$, and we recover in this way a result of Bloom and Shiffman [BS07, Example 3.5].

Our strategy for proving Theorem 1.4 consists in reducing to the univariate case. In §2, we consider the problem of studying the angle and radius discrepancies of an arbitrary 0-dimensional effective cycle Z in $(\mathbb{C}^\times)^n$ in terms of the angle and radius discrepancies of its direct images under all monomial projections of $(\mathbb{C}^\times)^n$ onto $\mathbb{C}^\times$. By applying a tomography process based on Fourier analysis, we show that the distribution of Z can be controlled in terms of the distribution of its projections (Theorem 2.3). In §3, we consider cycles defined by a system of Laurent polynomials with given support and we compute their direct image under monomial projections, in terms of sparse resultants. Theorem 1.4 then follows by applying the univariate Erdös-Turán's theorem combined with Theorem 2.3, and the basic properties of the sparse resultant.

In § 4, we study the asymptotic distribution of the roots of a sequence of systems of Laurent polynomials over $\mathbb{Z}$ and of random systems of Laurent polynomials over $\mathbb{C}$. In both situations, the key step consists in bounding from below the size of the relevant facet resultants. In the case of systems over $\mathbb{Z}$, this is trivial since these facet resultants are nonzero integer numbers. In the case of random systems over $\mathbb{C}$, the result follows from an estimate of the volume of a tube around an algebraic variety due to Beltrán and Pardo [BP07].

Acknowledgments. We thank Carlos Beltrán and Michael Shub for useful discussions and pointers to the literature. Part of this work was done while the authors met at Universitat de Barcelona and Université de Nice–Sophia Antipolis. We thank these institutions for their hospitality.

2. ANGLE AND RADIUS DISTRIBUTION IN THE MULTIVARIATE CASE

In this section, we show that the angle and radius discrepancies of an effective 0-dimensional cycle in the torus $(\mathbb{C}^\times)^n$ can be bounded in terms of the angle discrepancy of its image under monomial maps from $(\mathbb{C}^\times)^n$ to $\mathbb{C}^\times$.

Let Z be a nonzero effective 0-dimensional cycle of $(\mathbb{C}^\times)^n$, which we write as a finite sum

$$Z = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \boldsymbol{\xi}$$

with $m_{\boldsymbol{\xi}} \in \mathbb{N}$ and $\boldsymbol{\xi} \in (\mathbb{C}^\times)^n$. The *support* of Z is the finite subset of $(\mathbb{C}^\times)^n$ defined as $|Z| = \{\boldsymbol{\xi} \mid m_{\boldsymbol{\xi}} \geq 1\}$, and the *degree* of Z is the positive number $\deg(Z) = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}$.

Definition 2.1. Let $Z = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \boldsymbol{\xi}$ be a nonzero effective 0-dimensional cycle of $(\mathbb{C}^\times)^n$. For each $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n)$ and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_n)$ with $-\pi \leq \alpha_j < \beta_j \leq \pi$, $j = 1, \dots, n$, consider the cycle

$$Z_{\boldsymbol{\alpha},\boldsymbol{\beta}} = \sum_{-\alpha_j < \arg(\xi_j) \leq \beta_j} m_{\boldsymbol{\xi}} \boldsymbol{\xi}.$$

The *angle discrepancy* of Z is defined as

$$\Delta_{\mathrm{ang}}(Z) = \sup_{\boldsymbol{\alpha},\boldsymbol{\beta}} \left| \frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)} - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi} \right|.$$

Let $0 < \varepsilon < 1$ and consider also the cycle

$$Z_\varepsilon = \sum_{1-\varepsilon < |\xi_j| < (1-\varepsilon)^{-1}} m_{\boldsymbol{\xi}} \boldsymbol{\xi},$$

where ξ_j is the j-th coordinate of $\boldsymbol{\xi}$. The *radius discrepancy* of Z with respect to ε is defined as

$$\Delta_{\mathrm{rad}}(Z, \varepsilon) = 1 - \frac{\deg(Z_\varepsilon)}{\deg(Z)}.$$

We have $0 < \Delta_{\mathrm{ang}}(Z) \leq 1$. Observe also that $0 \leq \Delta_{\mathrm{rad}}(Z, \varepsilon) \leq 1$ and $\Delta_{\mathrm{rad}}(Z, \varepsilon) = 0$ for all ε if and only if $|Z| \subset (S^1)^n$.

For a lattice point $\boldsymbol{a} = (a_1, \dots, a_n) \in \mathbb{Z}^n$ we denote by $\chi^{\boldsymbol{a}} \colon (\mathbb{C}^\times)^n \to \mathbb{C}^\times$ the associated character, defined as $\chi^{\boldsymbol{a}}(\boldsymbol{\xi}) = \xi_1^{a_1} \dots \xi_n^{a_n}$ for $\boldsymbol{\xi} \in (\mathbb{C}^\times)^n$. The direct image of Z under $\chi^{\boldsymbol{a}}$ is the cycle of $\mathbb{C}^\times$ given by

$$\chi^{\boldsymbol{a}}_*(Z) = \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} \chi^{\boldsymbol{a}}(\boldsymbol{\xi}).$$

We also set

$$\theta(Z) = \sup_{\boldsymbol{a} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}} \frac{\Delta_{\mathrm{ang}}(\chi^{\boldsymbol{a}}_*(Z))}{\|\boldsymbol{a}\|_2^{\frac{1}{2}}}, \quad \rho(Z, \varepsilon) = \sum_{j=1}^{n} \Delta_{\mathrm{rad}}(\chi^{\boldsymbol{e}_j}_*(Z), \varepsilon), \tag{2.2}$$

where $\boldsymbol{e}_j$ denotes the j-th vector in the standard basis of $\mathbb{Z}^n$, and $\|\boldsymbol{a}\|_2$ is the Euclidean norm of the vector $\boldsymbol{a} \in \mathbb{Z}^n$. We have $0 < \theta(Z) \leq 1$ and $0 \leq \rho(Z, \varepsilon) \leq 1$.

Theorem 2.3. *Let Z be a nonzero effective 0-dimensional cycle of $(\mathbb{C}^\times)^n$. Then*

$$\Delta_{\mathrm{ang}}(Z) \leq 22n \left(\frac{8}{3}\right)^n (9 - \log(\theta(Z)))^{\frac{2}{3}(n-1)}\, \theta(Z)^{\frac{2}{3}}$$

and, for $0 < \varepsilon < 1$,

$$\Delta_{\mathrm{rad}}(Z, \varepsilon) \leq \rho(Z, \varepsilon).$$

The rest of this section is devoted to the proof of this result. For $\boldsymbol{u}, \boldsymbol{v} \in \mathbb{R}^n$ we write $\langle \boldsymbol{u}, \boldsymbol{v} \rangle = \sum_{j=1}^n u_j v_j$ for their standard inner product, and for $\boldsymbol{\xi} \in (\mathbb{C}^\times)^n$ we set $\arg(\boldsymbol{\xi}) = (\arg(\xi_1), \dots, \arg(\xi_n)) \in (-\pi, \pi]^n$.

Lemma 2.4. *Let Z be a nonzero effective 0-dimensional cycle of $(\mathbb{C}^\times)^n$ and $\boldsymbol{a} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$. Then*

$$\left| \frac{1}{\deg(Z)} \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\, \mathrm{e}^{\mathrm{i}\langle \boldsymbol{a}, \arg(\boldsymbol{\xi})\rangle} \right| \leq 2\pi \Delta_{\mathrm{ang}}(\chi^{\boldsymbol{a}}_*(Z)).$$

Note that for $\boldsymbol{a} = \boldsymbol{0}$ we get $\frac{1}{\deg(Z)} \sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\, \mathrm{e}^{\mathrm{i}\langle \boldsymbol{a}, \arg(\boldsymbol{\xi})\rangle} = 1$.

Proof. Set $D_0 = \#|Z|$ and $D = \deg(Z)$ for short. Let $-\pi \leq \nu_k < \pi$, $k = 1, \dots, D_0$, denote the inner products $\langle \boldsymbol{a}, \arg(\boldsymbol{\xi})\rangle$ modulo 2π for the different points $\boldsymbol{\xi}$ in the support of Z, and let m_k denote their corresponding multiplicity. We suppose that these numbers are arranged in increasing order, that is, $\nu_1 \leq \dots \leq \nu_{D_0}$.

For $-\pi < \nu \leq \pi$ set

$$N(\nu) = \sum_{\nu_k \leq \nu} m_k.$$

We have

$$\begin{aligned} \int_{-\pi}^{\pi} N(\nu)\, \mathrm{e}^{\mathrm{i}\nu}\, d\nu &= \sum_{k=1}^{D_0} \int_{\nu_k}^{\nu_{k+1}} \Big(\sum_{l \leq k} m_l\Big) \mathrm{e}^{\mathrm{i}\nu}\, d\nu \\ &= -\mathrm{i} \sum_k \Big(\sum_{l \leq k} m_l\Big)\, \mathrm{e}^{\mathrm{i}\nu} \Big|_{\nu_k}^{\nu_{k+1}} = \mathrm{i}\Big(D + \sum_k m_k\, \mathrm{e}^{\mathrm{i}\nu_k}\Big), \end{aligned} \tag{2.5}$$

where we have set $\nu_{D_0+1} = \pi$. On the other hand, an easy calculation shows that

$$\int_{-\pi}^{\pi} \frac{\nu + \pi}{2\pi}\, \mathrm{e}^{\mathrm{i}\nu}\, d\nu = \mathrm{i}. \tag{2.6}$$

Combining (2.5) and (2.6), we deduce that

$$\frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\, \mathrm{e}^{\mathrm{i}\langle \boldsymbol{a},\arg(\boldsymbol{\xi})\rangle} = \frac{1}{D}\sum_{k=1}^{D_0} m_k\, \mathrm{e}^{\mathrm{i}\nu_k} = \mathrm{i}\int_{-\pi}^{\pi}\Big(\frac{\nu+\pi}{2\pi}-\frac{N(\nu)}{D}\Big)\mathrm{e}^{\mathrm{i}\nu}\, d\nu.$$

Hence

$$\begin{aligned}\left|\frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\, \mathrm{e}^{\mathrm{i}\langle \boldsymbol{a},\arg(\boldsymbol{\xi})\rangle}\right| &\le \int_{-\pi}^{\pi}\left|\frac{\nu+\pi}{2\pi}-\frac{N(\nu)}{D}\right| d\nu \\ &= \int_{-\pi}^{\pi}\left|\frac{\nu+\pi}{2\pi}-\frac{\deg(\chi_*^{\boldsymbol{a}}(Z)_{0,\nu})}{\deg(\chi_*^{\boldsymbol{a}}(Z))}\right| d\nu \le 2\pi\Delta_{\mathrm{ang}}(\chi_*^{\boldsymbol{a}}(Z)),\end{aligned}$$

which concludes the proof □

Let $\alpha,\beta,\tau\in\mathbb{R}$ such that $\alpha\le\beta$ and $\tau>0$. We consider the function $h_{\alpha,\beta,\tau}\colon\mathbb{R}\to\mathbb{R}$ defined, for $x\in\mathbb{R}$, by

$$h_{\alpha,\beta,\tau}(x)=\begin{cases}0 & \text{if } x\le\alpha-\tau,\\ g\big(\frac{x-\alpha+\tau}{\tau}\big) & \text{if}\, \alpha-\tau<x\le\alpha,\\ 1 & \text{if } \alpha<x\le\beta,\\ g\big(\frac{\beta+\tau-x}{\tau}\big) & \text{if } \beta<x\le\beta+\tau,\\ 0 & \text{if } \beta+\tau<x,\end{cases}$$

with $g(x)=-2x^3+3x^2$. Lemma 2.7 below shows that $h_{\alpha,\beta,\tau}$ is an approximation of the characteristic function of the interval $[\alpha,\beta]$.

For $m\in\mathbb{N}$, we denote by $\mathcal{C}^m(\mathbb{R})$ the space of functions $f\colon\mathbb{R}\to\mathbb{R}$ having m continuous derivatives.

Lemma 2.7. *Let $\alpha,\beta,\tau\in\mathbb{R}$ such that $\alpha\le\beta$ and $\tau>0$. Then*

1. $h_{\alpha,\beta,\tau}\in\mathcal{C}^1(\mathbb{R})$;
2. $h_{\alpha,\beta,\tau}(x)=1$ *for* $x\in[\alpha,\beta]$, $h_{\alpha,\beta,\tau}(x)=0$ *for* $x\in(-\infty,\alpha-\tau]\cup[\beta+\tau,\infty)$, *and* $0\le h_{\alpha,\beta,\tau}(x)\le1$ *for all* $x\in\mathbb{R}$;
3. $\int_{-\infty}^{\infty}h_{\alpha,\beta,\tau}dx=\beta-\alpha+\tau$ *and, moreover,* $\int_{\alpha-\tau}^{\alpha}h_{\alpha,\beta,\tau}dx=\int_{\beta}^{\beta+\tau}h_{\alpha,\beta,\tau}dx=\frac{\tau}{2}$;
4. $\int_{-\infty}^{\infty}|h'_{\alpha,\beta,\tau}|dx=2$;
5. $h''_{\alpha,\beta,\tau}$ *is absolutely integrable and* $\int_{-\infty}^{\infty}|h''_{\alpha,\beta,\tau}|dx=\frac{6}{\tau}$.

Proof. By a direct calculation, we verify that the function g satisfies the following properties:

- $g(x)\ge0$ for all $x\in[0,1]$;
- $g(0)=g'(0)=0,\ g(1)=1,\ g'(1)=0$;
- $\int_0^1 g\,dx=\frac{1}{2}$;
- $\int_0^1|g'|\,dx=1$;
- $\int_0^1|g''|\,dx=3$.

The claim follows easily from these properties and the definition of $h_{\alpha,\beta,\tau}$. □

Suppose furthermore that $\beta-\alpha+2\tau<2\pi$. The support of $h_{\alpha,\beta,\tau}$ is then contained in an interval of length bounded by 2π, and so this function can be regarded as a function on $\mathbb{R}/2\pi\mathbb{Z}$. Consider its Fourier series

$$h_{\alpha,\beta,\tau}(x) \sim \sum_{a\in\mathbb{Z}} c_a \, \mathrm{e}^{\mathrm{i}ax},$$

with $c_a = \frac{1}{2\pi}\int_{-\pi}^{\pi} h_{\alpha,\beta,\tau}(x)\,\mathrm{e}^{-\mathrm{i}ax}\,dx$.

Lemma 2.8. *Let $\alpha \le \beta$ and $\tau > 0$ such that $\beta - \alpha + 2\tau < 2\pi$. Then $\sum_{a\in\mathbb{Z}} c_a \,\mathrm{e}^{\mathrm{i}ax}$ converges absolutely and uniformly on $\mathbb{R}/2\pi\mathbb{Z}$ to $h_{\alpha,\beta,\tau}$. Moreover, $c_0 = \frac{\beta-\alpha+\tau}{2\pi}$, and, for $a \neq 0$,*

$$|c_a| \le \min\Big\{\frac{1}{\pi\, a}, \frac{3}{\pi\,\tau\, a^2}\Big\}. \tag{2.9}$$

Proof. Lemma 2.7(1) implies that the series $\sum_{a\in\mathbb{Z}} c_a\,\mathrm{e}^{\mathrm{i}ax}$ converges absolutely and uniformly on $\mathbb{R}/2\pi\mathbb{Z}$ to the function $h_{\alpha,\beta,\tau}$. The computation of c_0 follows from Lemma 2.7(3). Integrating by parts, we deduce for $a \in \mathbb{Z}\setminus\{0\}$ that

$$c_a = \frac{1}{-2\pi\mathrm{i}a}\int_{-\pi}^{\pi} h'_{\alpha,\beta,\tau}(x)\,\mathrm{e}^{-\mathrm{i}ax}\,dx = \frac{1}{2\pi(-\mathrm{i}a)^2}\int_{-\pi}^{\pi} h''_{\alpha,\beta,\tau}(x)\,\mathrm{e}^{-\mathrm{i}ax}\,dx.$$

Hence, $|c_a| \le \frac{1}{2\pi a}\int_{-\pi}^{\pi}|h'_{\alpha,\beta,\tau}|dx$ and also $|c_a| \le \frac{1}{2\pi a^2}\int_{-\pi}^{\pi}|h''_{\alpha,\beta,\tau}|dx$. Then (2.9) follows by bounding these integrals with Lemma 2.7(4-5). □

Next, we apply Fourier analysis to control the angle discrepancy of Z in terms of the angle discrepancy of its direct image under monomial projections.

Lemma 2.10. *Let $n \ge 2$, $q \in \mathbb{Z}_{\ge 1}$, $\boldsymbol{\alpha} = (\alpha_1,\dots,\alpha_n)$ and $\boldsymbol{\beta} = (\beta_1,\dots,\beta_n)$ with $\alpha_j,\beta_j \in \mathbb{R}$ such that $-\pi \le \alpha_j < \beta_j < \pi$ and $\beta_j - \alpha_j + \frac{2}{q} < 2\pi$. Then*

$$\left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)} - \prod_{j=1}^{n}\frac{\beta_j-\alpha_j}{2\pi}\right| \le 2\,n\,\theta(Z) + \frac{3\,n}{2\,\pi\,q} + n\frac{2^{n+3}\sqrt{3}}{\pi^{n-1}}q^{\frac12}(9+\log(q))^{n-1}\,\theta(Z).$$

Proof. Set $\tau = \frac{1}{q}$ and $h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\boldsymbol{\nu}) = \prod_{j=1}^{n} h_{\alpha_j,\beta_j,\tau}(\nu_j)$ for $\boldsymbol{\nu} = (\nu_1,\dots,\nu_n) \in \mathbb{R}^n$. Set also $D = \deg(Z)$ and

$$\Sigma_1 = \left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{D} - \frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\arg(\boldsymbol{\xi}))\right|,$$

$$\Sigma_2 = \left|\frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}} h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\arg(\boldsymbol{\xi})) - \prod_{j=1}^{n}\frac{\beta_j-\alpha_j+\tau}{2\pi}\right|,$$

$$\Sigma_3 = \left|\prod_{j=1}^{n}\frac{\beta_j-\alpha_j+\tau}{2\pi} - \prod_{j=1}^{n}\frac{\beta_j-\alpha_j}{2\pi}\right|.$$

We will bound each of these quantities. For Σ_1, we consider the subset of $\mathbb{R}^n$ given by $I_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau} = \prod_{j=1}^{n}[\alpha_j-\tau,\beta_j+\tau]\setminus\prod_{j=1}^{n}[\alpha_j,\beta_j]$. Then

$$\Sigma_1 = \left|\frac{1}{D}\sum_{\arg(\boldsymbol{\xi})\in I_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}} m_{\boldsymbol{\xi}} h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\arg(\boldsymbol{\xi}))\right|.$$

For each $\boldsymbol{\xi}$ such that $\arg(\boldsymbol{\xi}) \in I_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}$, there is $1 \leq j \leq n$ such that either $\alpha_j - \tau < \arg(\xi_j) \leq \alpha_j$ or $\beta_j < \arg(\xi_j) < \beta_j + \tau$. Since $0 \leq h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\boldsymbol{\nu}) \leq 1$ for all $\boldsymbol{\nu}$, we have that Σ_1 is bounded above by

$$\frac{1}{D}\sum_{j=1}^{n}\big(\deg(\chi_*^{\boldsymbol{e}_j}(Z)_{\alpha_j-\tau,\alpha_j}) + \deg(\chi_*^{\boldsymbol{e}_j}(Z)_{\beta_j,\beta_j+\tau})\big).$$

By using the definition of Δ_{ang} and Lemma 2.4, we get

$$\frac{1}{D}\deg(\chi_*^{\boldsymbol{e}_j}(Z)_{\alpha_j-\tau,\alpha_j}) \leq \Delta_{\mathrm{ang}}(\chi_*^{\boldsymbol{e}_j}(Z)) + \frac{\tau}{2\pi} \leq \theta(Z) + \frac{\tau}{2\pi} = \theta(Z) + \frac{1}{2\pi q},$$

and a similar bound holds for $\deg(\chi_*^{\boldsymbol{e}_j}(Z)_{\beta_j,\beta_j+\tau})$. Hence,

$$\Sigma_1 \leq 2n\theta(Z) + \frac{n}{\pi q}. \tag{2.11}$$

Now we turn to Σ_2. Due to the conditions imposed on τ, we can regard $h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}$ as a function on $\mathbb{R}^n/2\pi\mathbb{Z}^n \simeq (-\pi,\pi]^n$. Let $\sum_{\boldsymbol{a}\in\mathbb{Z}^n} c_{\boldsymbol{a}}\,\mathrm{e}^{\mathrm{i}\langle\boldsymbol{a},\boldsymbol{\nu}\rangle}$ be its multivariate Fourier series. For $j=1,\dots,n$, we denote with $\sum_{a_j\in\mathbb{Z}} c_{j,a_j}\,\mathrm{e}^{\mathrm{i}a_j\nu_j}$ the Fourier series of $h_{\alpha_j,\beta_j,\tau}$. Then, for each $\boldsymbol{a}=(a_1,\dots,a_n)\in\mathbb{Z}^n$,

$$c_{\boldsymbol{a}} = \prod_{j=1}^{n} c_{j,a_j}.$$

In particular, $c_{\boldsymbol{0}} = \prod_{j=1}^{n}\frac{\beta_j-\alpha_j+\tau}{2\pi}$. The Fourier series of each $h_{\alpha_j,\beta_j,\tau}$ converges absolutely to this function, and so the same holds for the Fourier series of $h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}$. Hence,

$$h_{\boldsymbol{\alpha},\boldsymbol{\beta},\tau}(\boldsymbol{\nu}) = \sum_{\boldsymbol{a}\in\mathbb{Z}^n} c_{\boldsymbol{a}}\,\mathrm{e}^{\mathrm{i}\langle\boldsymbol{a},\boldsymbol{\nu}\rangle}$$

for $\boldsymbol{\nu}\in(-\pi,\pi]^n$. Applying Lemma 2.4, we obtain

$$\begin{aligned}\Sigma_2 &= \Big|\frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\sum_{\boldsymbol{a}\neq\boldsymbol{0}} c_{\boldsymbol{a}}\,\mathrm{e}^{\mathrm{i}\langle\boldsymbol{a},\arg(\boldsymbol{\xi})\rangle}\Big| \leq \sum_{\boldsymbol{a}\neq 0}|c_{\boldsymbol{a}}|\Big|\frac{1}{D}\sum_{\boldsymbol{\xi}} m_{\boldsymbol{\xi}}\,\mathrm{e}^{\mathrm{i}\langle\boldsymbol{a},\arg(\boldsymbol{\xi})\rangle}\Big| \\ &\leq 2\pi\,\theta(Z)\sum_{\boldsymbol{a}\neq 0}|c_{\boldsymbol{a}}|\|\boldsymbol{a}\|_2^{\frac{1}{2}} \leq 2^{n+1}\pi\,\theta(Z)\sum_{\boldsymbol{a}\geq\boldsymbol{0}}\Big(\sum_{s=1}^{n}\sqrt{a_s}\Big)|c_{\boldsymbol{a}}|\end{aligned} \tag{2.12}$$

For $s=1$,

$$\sum_{\boldsymbol{a}\geq\boldsymbol{0}}\sqrt{a_1}|c_{\boldsymbol{a}}| = \Big(\sum_{a_1\geq 0}\sqrt{a_1}|c_{1,a_1}|\Big)\prod_{j=2}^{n}\Big(\sum_{a_j\geq 0}|c_{j,a_j}|\Big). \tag{2.13}$$

Using the bounds in (2.9) we get, for $j = 2, \dots, n$,

$$\begin{aligned}\sum_{a_j \geq 0} |c_{j,a_j}| &\leq |c_{j,0}| + \sum_{a_j=1}^{3q} \frac{1}{a_j \pi} + \sum_{a_j=3q+1}^{\infty} \frac{3}{\pi\tau a_j^2} \\ &\leq \frac{\beta_j - \alpha_j + \tau}{2\pi} + \left(1 + \frac{1}{\pi}\int_1^{3q} \frac{dx}{x}\right) + \frac{3}{\pi\tau}\int_{3q}^{\infty} \frac{dx}{x^2} \\ &\leq \frac{2\pi}{2\pi} + 1 + \frac{1}{\pi}\log\left(\frac{3}{\tau}\right) + \frac{3}{\pi\tau}\frac{\tau}{3} \\ &\leq \frac{1}{\pi}\big(9 + \log q\big). \qquad (2.14)\end{aligned}$$

Similarly, we now bound

$$\begin{aligned}\sum_{a_1 \geq 0} \sqrt{a_1}|c_{1,a_1}| &\leq \sum_{a_1=1}^{3q} \frac{1}{\pi\sqrt{a_1}} + \sum_{a_1=3q+1}^{\infty} \frac{3}{\pi\tau\, a_1\sqrt{a_1}} \\ &\leq \frac{1}{\pi}\int_0^{3q} x^{-\frac{1}{2}}dx + \frac{3}{\pi\tau}\int_{3q}^{\infty} x^{-\frac{3}{2}}dx \\ &\leq \frac{2}{\pi}\left(\frac{\tau}{3}\right)^{-\frac{1}{2}} + \frac{6}{\pi\tau}\left(\frac{\tau}{3}\right)^{\frac{1}{2}} \\ &\leq \frac{4\sqrt{3}}{\pi} q^{\frac{1}{2}}. \qquad (2.15)\end{aligned}$$

It follows from (2.12), (2.14) and (2.15) that

$$\Sigma_2 \leq n\frac{2^{n+3}\sqrt{3}}{\pi^{n-1}} q^{\frac{1}{2}}(9 + \log(q))^{n-1}\,\theta(Z). \qquad (2.16)$$

Next we consider Σ_3. Set $\phi(t) = \prod_{j=1}^{n} \frac{\beta_j - \alpha_j + \tau t}{2\pi}$ for $t \in \mathbb{R}$. There exists $0 < t_0 < 1$ such that $\Sigma_3 = |\phi(1) - \phi(0)| = |\phi'(t_0)|$ and so

$$\Sigma_3 \leq \sup_{0<t_0<1} |\phi'(t_0)| \leq \sum_{j=1}^{n} \frac{\tau}{2\pi} \prod_{\ell \neq j} \frac{\beta_\ell - \alpha_\ell + \tau}{2\pi} \leq n\tau\frac{(2\pi)^{n-1}}{(2\pi)^n} = \frac{n}{2\pi q}.$$

Finally, we collect (2.11), (2.16) and the above inequality to get

$$\begin{aligned}\left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)} - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi}\right| &\leq \Sigma_1 + \Sigma_2 + \Sigma_3 \\ &\leq 2n\theta(Z) + \frac{n}{\pi q} + n\frac{2^{n+3}\sqrt{3}}{\pi^{n-1}} q^{\frac{1}{2}}(9 + \log(q))^{n-1}\,\theta(Z) + \frac{n}{2\pi q} \\ &= 2n\theta(Z) + \frac{3}{2}\frac{n}{\pi q} + n\frac{2^{n+3}\sqrt{3}}{\pi^{n-1}} q^{\frac{1}{2}}(9 + \log(q))^{n-1}\,\theta(Z).\end{aligned}$$

□

Proof of Theorem 2.3. For the radius discrepancy, we have

$$|Z_\varepsilon| = \bigcap_{j=1}^{n} \left\{\boldsymbol{\xi} \in |Z| \;\middle|\; 1 - \varepsilon < |\xi_j| < \frac{1}{1-\varepsilon}\right\}.$$

By taking complements in this equality and considering the corresponding multiplicities, we deduce that

$$\deg(Z) - \deg(Z_\varepsilon) \leq \sum_{j=1}^{n} \deg(\chi_*^{\boldsymbol{e}_j}(Z)) - \deg(\chi_*^{\boldsymbol{e}_j}(Z)_\varepsilon).$$

Hence, $\Delta_{\mathrm{rad}}(Z,\varepsilon) \leq \sum_{j=1}^{n} \Delta_{\mathrm{rad}}\big(\chi_*^{\boldsymbol{e}_j}(Z,\varepsilon)\big) = \rho(Z,\varepsilon)$, as stated.

We now consider the bound for the angle discrepancy. For $n=1$, $\Delta_{\mathrm{ang}}(Z) \leq \theta(Z)$ and so the bound in the claim is trivial. Hence, we suppose that $n \geq 2$. Put then $\zeta(Z) = (9-\log(\theta(Z)))^{\frac{2}{3}(n-1)}\,\theta(Z)^{2/3} \in \mathbb{R}_{>0}$ for short and set

$$q = \left\lfloor \frac{9^{\frac{2}{3}(n-1)}}{\zeta(Z)} \right\rfloor .$$

Suppose also that $q \geq 1$. Then

$$\frac{9^{\frac{2}{3}(n-1)}}{2\zeta(Z)} < q \leq \frac{9^{\frac{2}{3}(n-1)}}{\zeta(Z)} \leq \frac{1}{\theta(Z)}. \tag{2.17}$$

Let $\boldsymbol{\alpha} = (\alpha_1,\dots,\alpha_n)$ and $\boldsymbol{\beta} = (\beta_1,\dots,\beta_n)$ with $-\pi \leq \alpha_j < \beta_j \leq \pi$. Consider first the case when $\beta_j - \alpha_j \leq \pi$. In particular, $\beta_j - \alpha_j + \frac{2}{q} < 2\pi$. Applying Lemma 2.10 and the inequalities (2.17), we deduce that the quantity $\left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)} - \prod_{j=1}^{n} \frac{\beta_j-\alpha_j}{2\pi}\right|$ is bounded above by

$$2n\frac{\zeta(Z)}{9^{\frac{2}{3}(n-1)}} + \frac{3\,n}{2\,\pi}\frac{2\,\zeta(Z)}{9^{\frac{2}{3}(n-1)}} + n\frac{2^{n+3}\sqrt{3}}{\pi^{n-1}}\left(\frac{9^{\frac{2}{3}(n-1)}}{\zeta(Z)}\right)^{\frac{1}{2}}(9-\log(\theta(Z)))^{n-1}\,\theta(Z).$$

Since $n \geq 2$, this quantity can be bounded by

$$\begin{aligned} n\bigg(\frac{2}{9^{\frac{2}{3}(n-1)}} + \frac{3}{9^{\frac{2}{3}(n-1)}\pi} + \frac{2^{n+3}9^{\frac{1}{3}(n-1)}\sqrt{3}}{\pi^{n-1}}\bigg)\zeta(Z)& \\ \leq n\bigg(1 + \frac{2^3\sqrt{3}\pi}{9^{\frac{1}{3}}}\bigg(\frac{2\cdot 9^{\frac{1}{3}}}{\pi}\bigg)^{n}\bigg)\zeta(Z) &\leq 22\,n\left(\frac{4}{3}\right)^{n}\zeta(Z), \end{aligned}$$

as it can be easily verified that $\frac{2\cdot 9^{\frac{1}{3}}}{\pi} < \frac{4}{3}$ and $\frac{2^3\sqrt{3}\pi}{9^{\frac{1}{3}}} < 21$.

If $q=0$, then $\frac{9^{\frac{2}{3}(n-1)}}{\zeta(Z)} < 1$, which implies that $\Delta_{\mathrm{ang}}(Z) \leq 1 \leq \zeta(Z)$. Hence, in the case when $-\pi \leq \alpha_j < \beta_j \leq \pi$ for all j, it holds

$$\left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)} - \prod_{j=1}^{n}\frac{\beta_j-\alpha_j}{2\pi}\right| \leq 22\,n\left(\frac{4}{3}\right)^{n}\zeta(Z). \tag{2.18}$$

Now, if $\beta_j - \alpha_j > \pi$ for a j, we subdivide each of those intervals $(\alpha_j,\beta_j]$ into two subintervals of length $\leq \pi$. We can then decompose $Z_{\boldsymbol{\alpha},\boldsymbol{\beta}}$ as the sum of at most 2^n cycles of the form $Z_{\boldsymbol{\nu}_{0,\sigma},\boldsymbol{\nu}_{1,\sigma}}$ where the jth coordinate of $\boldsymbol{\nu}_{0,\sigma}$ (respectively $\boldsymbol{\nu}_{1,\sigma}$) is either α_j (respectively $\frac{\alpha_j+\beta_j}{2}$) or $\frac{\alpha_j+\beta_j}{2}$ (respectively β_j). Also, we can expand the product

$$\prod_{j=1}^{n}\frac{\beta_j-\alpha_j}{2\pi}$$

as the sum of the volumes of sets $\prod_{j=1}^{m}(\frac{\nu_{0,\sigma,j}}{2\pi},\frac{\boldsymbol{\nu}_{1,\sigma,j}}{2\pi}]$. From here, we easily get that

$$\left|\frac{\deg(Z_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{\deg(Z)}-\prod_{j=1}^{n}\frac{\beta_j-\alpha_j}{2\pi}\right|\leq\sum_{\sigma}\left|\frac{\deg(Z_{\boldsymbol{\nu}_{0,\sigma},\boldsymbol{\nu}_{1,\sigma}})}{\deg(Z)}-\prod_{j=1}^{n}\frac{\nu_{1,\sigma,j}-\nu_{0,\sigma,j}}{2\pi}\right|.$$

and the claim follows applying the bound (2.18), which has to be multiplied by 2^n. Altogether, we get

$$\Delta_{\mathrm{ang}}(Z)\leq 22\,n\left(\frac{8}{3}\right)^{n}\zeta(Z),$$

which concludes the proof. □

3. Bounds for the discrepancy in terms of sparse resultants

In this section, we consider cycles defined by a system of Laurent polynomials with given support. We first compute their direct image under monomial projections, in terms of sparse resultants. Then, we will derive Theorem 1.4 from the univariate Erdös-Turán's theorem and the results in the previous section.

We recall the definition of the sparse resultant, as presented in [DS12]. This definition differs from the classical one which can be found, for instance, in [GKZ94, CLO05, PS93].

Let $M\simeq\mathbb{Z}^n$ be a lattice of rank n and $N=M^{\vee}=\operatorname{Hom}(M,\mathbb{Z})$ its dual lattice. Set $M_{\mathbb{R}}=M\otimes\mathbb{R}$ and $N_{\mathbb{R}}=N\otimes\mathbb{R}$. The pairing between $b\in M_{\mathbb{R}}$ and $v\in N_{\mathbb{R}}$ will be denoted by $\langle b,v\rangle$. Let $\mathcal{A}_0,\dots,\mathcal{A}_n$ be a family of $n+1$ non-empty finite subsets of M and put $\boldsymbol{\mathcal{A}}=(\mathcal{A}_0,\dots,\mathcal{A}_n)$. Let $\boldsymbol{u}_i=\{u_{i,\boldsymbol{a}}\}_{a\in\mathcal{A}_i}$, $i=0,\dots,n$, be a group of $\#\mathcal{A}_i$ variables and set $\boldsymbol{u}=\{\boldsymbol{u}_0,\dots,\boldsymbol{u}_n\}$. For each i, let F_i be the general polynomial with support $\mathcal{A}_i$, defined as

$$F_i=\sum_{\boldsymbol{a}\in\mathcal{A}_i}u_{i,\boldsymbol{a}}\boldsymbol{x}^{\boldsymbol{a}}\in\mathbb{C}[\boldsymbol{u}][M],\tag{3.1}$$

where $\mathbb{C}[\boldsymbol{u}][M]\simeq\mathbb{C}[\boldsymbol{u}][x_1^{\pm1},\dots,x_n^{\pm1}]$ is the group algebra.

Let $\mathbb{T}_M=\operatorname{Hom}(M,\mathbb{C}^{\times})\simeq(\mathbb{C}^{\times})^n$ be the algebraic torus associated to M, and set

$$W_{\boldsymbol{\mathcal{A}}}=\Big\{(\boldsymbol{x},\boldsymbol{u})\in\mathbb{T}_M\times\prod_{i=0}^{n}\mathbb{P}(\mathbb{C}^{\mathcal{A}_i})\;\Big|\;F_i(\boldsymbol{u}_i,\boldsymbol{x})=0\Big\}.$$

This is an irreducible subvariety defined over $\mathbb{Q}$, of codimension $n+1$. Let

$$\pi\colon\mathbb{T}_M\times\prod_{i=0}^{n}\mathbb{P}(\mathbb{C}^{\mathcal{A}_i})\longrightarrow\prod_{i=0}^{n}\mathbb{P}(\mathbb{C}^{\mathcal{A}_i})$$

be the projection onto the second factor. The direct image $W_{\boldsymbol{\mathcal{A}}}$ under this map is the Weil divisor of $\prod_{i=0}^{n}\mathbb{P}(\mathbb{C}^{\mathcal{A}_i})$ given by

$$\pi_*(W_{\mathcal{A}})=\begin{cases}\deg(\pi|_{W_{\boldsymbol{\mathcal{A}}}})\overline{\pi(W_{\boldsymbol{\mathcal{A}}})} & \text{if } \operatorname{codim}(\overline{\pi(W_{\boldsymbol{\mathcal{A}}})})=1,\\ 0 & \text{if } \operatorname{codim}(\overline{\pi(W_{\boldsymbol{\mathcal{A}}})})\geq 2.\end{cases}$$

The *sparse resultant* associated to $\boldsymbol{\mathcal{A}}$, denoted $\operatorname{Res}_{\boldsymbol{\mathcal{A}}}$, is defined as any primitive equation in $\mathbb{Z}[\boldsymbol{u}]$ of this Weil divisor. It is well-defined up to a sign. When $\pi_*(W_{\mathcal{A}})=0$, we set $\operatorname{Res}_{\boldsymbol{\mathcal{A}}}=1$ for convenience.

According to the definition above, sparse resultants are not necessarily irreducible. If, as in [DS12], we denote by $\operatorname{Elim}_{\boldsymbol{\mathcal{A}}}$ what is classically called the sparse resultant (for instance in [GKZ94, CLO05, PS93]), then $\mathrm{Res}_{\boldsymbol{\mathcal{A}}} \neq 1$ if and only if $\mathrm{Elim}_{\boldsymbol{\mathcal{A}}} \neq 1$ and, if this is the case,

$$\mathrm{Res}_{\boldsymbol{\mathcal{A}}} = \pm \, \mathrm{Elim}_{\boldsymbol{\mathcal{A}}}^{\deg(\pi|_{W_{\boldsymbol{\mathcal{A}}}})}.$$

We recall the basic properties of the sparse resultant that we will need in the sequel. For simplicity, we will restrict to the case when $M = N = \mathbb{Z}^n$, so that each $\mathcal{A}_i$ is a finite subset of $\mathbb{Z}^n$. Consider the convex hull $Q_i = \mathrm{conv}(\mathcal{A}_i)$, which is a lattice polytope of $\mathbb{R}^n$. The *mixed volume* of convex bodies $P_1, \dots, P_n$ of $\mathbb{R}^n$ is defined as

$$\mathrm{MV}(P_1, \dots, P_n) = \sum_{j=1}^{n} (-1)^{n-j} \sum_{1 \le i_1 < \cdots < i_j \le n} \mathrm{vol}(P_{i_1} + \cdots + P_{i_j}).$$

Write $\mathbb{C}[\boldsymbol{x}^{\pm 1}] = \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ for short. The *height* of a Laurent polynomial $f = \sum_{\boldsymbol{a} \in \mathbb{Z}^n} \alpha_{\boldsymbol{a}} \boldsymbol{x}^{\boldsymbol{a}} \in \mathbb{C}[\boldsymbol{x}^{\pm 1}]$ is defined as

$$\mathrm{h}(f) = \log \big(\max_{\boldsymbol{a}} |\alpha_{\boldsymbol{a}}| \big).$$

Proposition 3.2. *Let notations be as above. Then, the sparse resultant is a multihomogeneous polynomial of partial degrees*

$$\deg_{\boldsymbol{u}_i}(\mathrm{Res}_{\boldsymbol{\mathcal{A}}}) = \mathrm{MV}(Q_0, \dots, Q_{i-1}, Q_{i+1}, \dots, Q_n)$$

and height bounded by

$$\mathrm{h}(\mathrm{Res}_{\boldsymbol{\mathcal{A}}}) \le \sum_{i=0}^{n} \mathrm{MV}(Q_0, \dots, Q_{i-1}, Q_{i+1}, \dots, Q_n) \log(\#\mathcal{A}_i).$$

Proof. The formula for the partial degrees is classical, see for instance [GKZ94]. In the present form, it can be found in [DS12]. The bound for the height is given by [Som04, Theorem 1] for the case when the resultant depends on all the groups of variables $\boldsymbol{u}_0, \dots, \boldsymbol{u}_n$. The general case can be found in [DS12]. □

For a family of Laurent polynomials $f_i \in \mathbb{C}[\boldsymbol{x}^{\pm 1}]$ with $\mathrm{supp}(f_i) \subset \mathcal{A}_i$, $i = 0, \dots, n$, we write

$$\mathrm{Res}_{\boldsymbol{\mathcal{A}}}(f_0, \dots, f_n)$$

for the evaluation of the resultant at their coefficients. The following is the multiplicativity formula for sparse resultants.

Proposition 3.3. *Let $\boldsymbol{\mathcal{A}} = (\mathcal{A}_0, \dots, \mathcal{A}_n)$ and $\boldsymbol{\mathcal{A}}' = (\mathcal{A}_0', \dots, \mathcal{A}_n')$ be two families of non-empty finite subsets of $\mathbb{Z}^n$ and set $\boldsymbol{\mathcal{A}} + \boldsymbol{\mathcal{A}}' = (\mathcal{A}_0 + \mathcal{A}_0', \dots, \mathcal{A}_n + \mathcal{A}_n')$. Let $f_i, f_i' \in \mathbb{C}[\boldsymbol{x}^{\pm 1}]$, $i = 0, \dots, n$, be Laurent polynomials with support contained in $\mathcal{A}_i$ and in $\mathcal{A}_i'$, respectively. Then*

$$\mathrm{Res}_{\boldsymbol{\mathcal{A}}+\boldsymbol{\mathcal{A}}'}(f_0 f_0', \dots, f_n f_n') = \mathrm{Res}_{\boldsymbol{\mathcal{A}}}(f_0, \dots, f_n) \cdot \mathrm{Res}_{\boldsymbol{\mathcal{A}}'}(f_0', \dots, f_n').$$

Proof. The case when the resultant depends on all the groups of variables $\boldsymbol{u}_0, \dots, \boldsymbol{u}_n$ is done in [PS93, Proposition 7.1]. The general case can be found in [DS12]. □

The *support function* of a compact subset $P \subset \mathbb{R}^n$ is the function $h_P \colon \mathbb{R}^n \to \mathbb{R}$ defined, for $\boldsymbol{v} \in \mathbb{R}^n$, as

$$h_P(\boldsymbol{v}) = \inf_{\boldsymbol{a} \in P} \langle \boldsymbol{a}, \boldsymbol{v} \rangle.$$

Let $\mathcal{B}$ be a finite subset of $\mathbb{Z}^n$ and $f = \sum_{\boldsymbol{b}\in\mathcal{B}} \beta_{\boldsymbol{b}} x^{\boldsymbol{b}}$ a Laurent polynomial with support contained in $\mathcal{B}$. For $\boldsymbol{v} \in \mathbb{R}^n$, we set

$$\mathcal{B}^{\boldsymbol{v}} = \{\boldsymbol{b} \in \mathcal{B} \mid \langle \boldsymbol{b}, \boldsymbol{v}\rangle = h_{\mathrm{conv}(\mathcal{B})}(\boldsymbol{v})\}, \quad f^{\boldsymbol{v}} = \sum_{\boldsymbol{b}\in\mathcal{B}^{\boldsymbol{v}}} \beta_{\boldsymbol{b}} x^{\boldsymbol{b}}.$$

A vector $\boldsymbol{v} \in \mathbb{Z}^n$ is *primitive* if the gcd of its coordinates is equal to 1.

Definition 3.4. Let $\mathcal{A}_1, \dots, \mathcal{A}_n \subset \mathbb{Z}^n$ be a family of n non-empty finite subsets, $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{0\}$, and $\boldsymbol{v}^\perp \subset \mathbb{R}^n$ the orthogonal subspace. Then $\mathbb{Z}^n \cap \boldsymbol{v}^\perp$ is a lattice of rank $n-1$ and, for $i = 1, \dots, n$, there exists $\boldsymbol{b}_{i,\boldsymbol{v}} \in \mathbb{Z}^n$ such that $\mathcal{A}_i^{\boldsymbol{v}} - \boldsymbol{b}_{i,\boldsymbol{v}} \subset \mathbb{Z}^n \cap \boldsymbol{v}^\perp$. The *facet resultant* of $\mathcal{A}_1, \dots, \mathcal{A}_n$ in the direction of $\boldsymbol{v}$, denoted $\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}$, is defined as the resultant of the family of finite subsets $\mathcal{A}_i^{\boldsymbol{v}} - \boldsymbol{b}_{i,\boldsymbol{v}} \subset \mathbb{Z}^n \cap \boldsymbol{v}^\perp$.

Let $f_i \in \mathbb{C}[x_1^{\pm1}, \dots, x_n^{\pm1}]$, $i = 1, \dots, n$, with $\mathrm{supp}(f_i) \subset \mathcal{A}_i$. For each i, write $f_i^{\boldsymbol{v}} = \boldsymbol{x}^{\boldsymbol{b}_{i,\boldsymbol{v}}} g_{i,\boldsymbol{v}}$ for a Laurent polynomial $g_{i,\boldsymbol{v}} \in \mathbb{C}[\mathbb{Z}^n \cap \boldsymbol{v}^\perp] \simeq \mathbb{C}[y_1^{\pm1}, \dots, y_{n-1}^{\pm1}]$ with $\mathrm{supp}(g_{i,\boldsymbol{v}}) \subset \mathcal{A}_i^{\boldsymbol{v}} - \boldsymbol{b}_{i,\boldsymbol{v}}$. The expression

$$\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}})$$

is defined as the evaluation of this facet resultant at the coefficients of the $g_{i,\boldsymbol{v}}$. These constructions are independent of the choice of the vectors $\boldsymbol{b}_{i,\boldsymbol{v}}$.

We have that $\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}} \neq 1$ only if $\boldsymbol{v}$ is an inner normal to a facet of the Minkowski sum $\sum_{i=1}^n Q_i$. In particular, the number of non-trivial facet resultants of the family $\mathcal{A}_1, \dots, \mathcal{A}_n$ is finite.

With notations as in Definition 3.4, write $\boldsymbol{f} = (f_1, \dots, f_n)$ for short. We denote by $V(\boldsymbol{f})_0 \subset \left(\mathbb{C}^\times\right)^n$ the set of isolated solutions in the torus of the system of equations $f_1 = \dots = f_n = 0$ and we set

$$Z(\boldsymbol{f}) = \sum_{\boldsymbol{\xi}\in V(\boldsymbol{f})_0} \mathrm{mult}(\boldsymbol{\xi}|\boldsymbol{f})\boldsymbol{\xi}$$

for the associated 0-dimensional cycle, where $\mathrm{mult}(\boldsymbol{\xi}|\boldsymbol{f})$ denotes the intersection multiplicity of $\boldsymbol{f}$ at a point $\boldsymbol{\xi}$. For $f_0 \in \mathbb{C}[\boldsymbol{x}^{\pm1}]$, we set

$$f_0(Z(\boldsymbol{f})) = \prod_{\boldsymbol{\xi}} f_0(\boldsymbol{\xi})^{\mathrm{mult}(\boldsymbol{\xi}|\boldsymbol{f})}.$$

The following result is known as the Poisson formula for sparse resultants.

Proposition 3.5. *Let $\boldsymbol{\mathcal{A}} = (\mathcal{A}_0, \dots, \mathcal{A}_n)$ be a family of non-empty finite subsets of $\mathbb{Z}^n$ and $f_i \in \mathbb{C}[\boldsymbol{x}^{\pm1}]$ a Laurent polynomial with* $\mathrm{supp}(f_i) \subset \mathcal{A}_i$*, $i = 0, \dots, n$. Suppose that* $\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}}) \neq 0$ *for all $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$. Then*

$$\mathrm{Res}_{\boldsymbol{\mathcal{A}}}(f_0, f_1, \dots, f_n) = \pm\left(\prod_{\boldsymbol{v}} \mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}})^{-h_{\mathcal{A}_0}(\boldsymbol{v})}\right) f_0(Z(f_1, \dots, f_n)),$$

the product being over all primitive elements $\boldsymbol{v} \in \mathbb{Z}^n$.

Proof. The case when the resultant depends on all the groups of variables $\boldsymbol{u}_0, \dots, \boldsymbol{u}_n$ can be derived from [PS93] (see also [Min03]). The general case can be found in [DS12]. □

From now on, we fix a family of non-empty finite subsets $\mathcal{A}_1, \dots, \mathcal{A}_n$ of $\mathbb{Z}^n$ such that $\mathrm{MV}(Q_1, \dots, Q_n) \geq 1$, with $Q_i = \mathrm{conv}(\mathcal{A}_i)$. We consider also a family of Laurent polynomials $f_1, \dots, f_n \in \mathbb{C}[\boldsymbol{x}^{\pm1}]$ with $\mathrm{supp}(f_i) \subset \mathcal{A}_i$, such that $\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}}) \neq 0$

for all $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{0\}$. By Bernstein-Kushnirenko-Khovanskii's theorem [Ber75, Theorem B],

$$\deg(Z(\boldsymbol{f})) = \mathrm{MV}(Q_1, \dots, Q_n) \geq 1.$$

Consider the projection $\pi_{\boldsymbol{e}_1} \colon \mathbb{R}^n \to \mathbb{R}^{n-1}$ given by $\pi_{\boldsymbol{e}_1}(x_1, x_2, \dots, x_n) = (x_2, \dots, x_n)$. If we regard each f_i as a Laurent polynomial in the group of variables $\boldsymbol{x}' := (x_2, \dots, x_n)$ with coefficients in the ring $\mathbb{C}[x_1{}^{\pm 1}]$, its support with respect to $\boldsymbol{x}'$ is contained in the finite subset $\pi_{\boldsymbol{e}_1}(\mathcal{A}_i)$ of $\mathbb{R}^{n-1}$. Set then

$$R(\boldsymbol{f}) = \mathrm{Res}_{\pi_{e_1}(\mathcal{A}_1), \dots, \pi_{e_1}(\mathcal{A}_n)} \big(f_1(x_1, \boldsymbol{x}'), \dots, f_n(x_1, \boldsymbol{x}')\big) \in \mathbb{C}[x_1^{\pm 1}]$$

for the evaluation of the resultant at these coefficients in $\mathbb{C}[x_1^{\pm 1}]$.

Recall that the sup-norm of a Laurent polynomial $f \in \mathbb{C}[\boldsymbol{x}^{\pm 1}]$ is defined as $\|f\|_{\sup} = \sup_{\boldsymbol{w} \in (S^1)^n} |f(\boldsymbol{w})|$. In general, it holds that

$$\mathrm{h}(f) \leq \log \|f\|_{\sup} \tag{3.6}$$

This is a consequence of Cauchy's formula for the coefficients of the Taylor expansion of a holomorphic function on $\mathbb{C}^N$ (see for instance [Som04, page 1255]). For a hyperplane $H \subset R^n$, we denote by MV_H the mixed volume associated to the Euclidean volume on H, defined for families of $n-1$ convex bodies of H.

The following result gives a bound for the sup-norm of $R(\boldsymbol{f})$. Its proof is a variant of that for [Som04, Lemma 1.3].

Lemma 3.7. *Let notations be as above. Then*

$$\log \|R(\boldsymbol{f})\|_{\sup} \leq \sum_{j=1}^{n} \mathrm{MV}_{\boldsymbol{e}_1^{\perp}}(\{\pi_{\boldsymbol{e}_1}(Q_\ell)\}_{\ell \neq j}) \log \|f_j\|_{\sup}.$$

Proof. Let $k \geq 1$. Proposition 3.3 implies that

$$R(\boldsymbol{f})^{k^n} = \mathrm{Res}_{k\pi_{\boldsymbol{e}_1}(\mathcal{A}_1), \dots, k\pi_{\boldsymbol{e}_1}(\mathcal{A}_n)}(f_1(x_1, \boldsymbol{x}')^k, \dots, f_n(x_1, \boldsymbol{x}')^k), \tag{3.8}$$

where $k\pi_{\boldsymbol{e}_1}(\mathcal{A}_j)$ denotes the pointwise sum of k copies of $\pi_{\boldsymbol{e}_1}(\mathcal{A}_j)$. For short, set $R_k = \mathrm{Res}_{k\pi_{\boldsymbol{e}_1}(\mathcal{A}_1), \dots, k\pi_{\boldsymbol{e}_1}(\mathcal{A}_n)}$ and $\boldsymbol{f}^k = (f_1^k, \dots, f_n^k)$, so that the identity above can be rewritten as $R(\boldsymbol{f})^{k^n} = R_k(\boldsymbol{f}^k)$. By Proposition 3.2, the partial degrees of this resultant are given by

$$\deg_{\boldsymbol{u}_j}(R_k) = \mathrm{MV}_{\boldsymbol{e}_1^{\perp}}(\{k\pi_{\boldsymbol{e}_1}(Q_\ell)\}_{\ell \neq j}) = k^{n-1} \, \mathrm{MV}_{\boldsymbol{e}_1^{\perp}}(\{\pi_{\boldsymbol{e}_1}(Q_\ell)\}_{\ell \neq j}),$$

where $\boldsymbol{u}_j$ is a group of $\# k\mathcal{A}_j$ variables, for $j = 1, \dots, n$. In particular, the logarithm of its number of monomials is bounded above as

$$\begin{aligned} \log(\#\,\mathrm{supp}(R_k)) &\leq \log\bigg(\prod_{j=1}^{n} \binom{\# k\mathcal{A}_j + \deg_{\boldsymbol{u}_j}(R_k)}{\# k\mathcal{A}_j}\bigg) \\ &\leq \sum_{j=1}^{n} \deg_{\boldsymbol{u}_j}(R_k) \log(\# k\mathcal{A}_j + 1) = O(k^{n-1} \log(k+1)), \end{aligned}$$

since $\# k\mathcal{A}_j \leq (k+1)^{c_1 n}$ for a constant c_1 independent of k. By Proposition 3.2, the height of this resultant is bounded above by

$$\mathrm{h}(R_k) \leq \sum_{j=1}^{n} \deg_{\boldsymbol{u}_j}(R_k) \log(\# k\mathcal{A}_j) = O(k^{n-1} \log(k+1)).$$

Let $w_1 \in S^1$. Using (3.8), (3.6), the previous bounds, and the fact that $\|f_j^k\|_{\sup} = \|f_j\|_{\sup}^k$, we deduce that

$$\begin{aligned} k^n \log \|R(\boldsymbol{f})\|_{\sup} &= \log \|R_k(\boldsymbol{f}^k)\|_{\sup} \\ &\le \mathrm{h}(R_k) + \sum_{j=1}^{n} \deg_{\boldsymbol{u}_j}(R_j) \log \|f_j^k\|_{\sup} + \log(\#\operatorname{supp}(R_k)) \\ &= k^n \Big(\sum_{j=1}^{n} \mathrm{MV}_{\boldsymbol{e}_1^\perp}(\{\pi_{\boldsymbol{e}_1}(Q_\ell)\}_{\ell \neq j}) \log \|f_j\|_{\sup} \Big) + O(k^{n-1} \log(k+1)). \end{aligned}$$

The result then follows by diving both members of this inequality by k^n and letting $k \to \infty$. □

For $\boldsymbol{a} \in \mathbb{Z}^n \setminus \{0\}$ and z an additional variable, set

$$E_{\boldsymbol{a}}(\boldsymbol{f}) = \mathrm{Res}_{\{\boldsymbol{0},\boldsymbol{a}\},\mathcal{A}_1,\dots,\mathcal{A}_n}(z - \boldsymbol{x}^{\boldsymbol{a}}, f_1, \dots, f_n) \in \mathbb{C}[z].$$

Due to the Poisson formula for sparse resultants given in Proposition 3.5, we have that $Z(E_{\boldsymbol{a}}(\boldsymbol{f})) = \chi_*^{\boldsymbol{a}}(Z(\boldsymbol{f}))$, and so $E_{\boldsymbol{a}}(\boldsymbol{f})$ can be regarded as an elimination polynomial for the cycle $Z(\boldsymbol{f})$ with respect to the monomial projection $\chi^{\boldsymbol{a}}$.

By [DS12], there exists $m \in \mathbb{Z}$ such that $E_{\boldsymbol{e}_1}(\boldsymbol{f})(x_1) = x_1^m R(\boldsymbol{f})$ (see also [CLO05, Proposition 5.15, Chapter 3] for a proof for the resultant of dense polynomials). Hence, Lemma 3.7 can be regarded as a bound for the sup-norm of the elimination polynomial $E_{\boldsymbol{e}_1}(\boldsymbol{f})$. Our next step is to extend this result to an arbitrary $\boldsymbol{a}$. Recall that $\pi_{\boldsymbol{a}} \colon \mathbb{R}^n \to \boldsymbol{a}^\perp$ denotes the orthogonal projection onto the hyperplane $\boldsymbol{a}^\perp \subset \mathbb{R}^n$.

Lemma 3.9. *Let notation be as above. Then*

$$\log \|E_{\boldsymbol{a}}(\boldsymbol{f})\|_{\sup} \le \|\boldsymbol{a}\|_2 \sum_{j=1}^{n} \mathrm{MV}_{\boldsymbol{a}^\perp}(\{\pi_{\boldsymbol{a}}(Q_\ell)\}_{\ell \neq j}) \log \|f_j\|_{\sup}.$$

Proof. Consider first the case when $\boldsymbol{a} \in \mathbb{Z}^n$ is primitive. The quotient $\mathbb{Z}^n/\boldsymbol{a}\mathbb{Z}$ is torsion-free and so $\boldsymbol{a}$ can be completed to a basis of $\mathbb{Z}^n$. Equivalently, there is an invertible matrix $A \in \mathrm{SL}_n(\mathbb{Z})$ with first row $\boldsymbol{a}$. Set $\boldsymbol{a}, \boldsymbol{a}_2, \dots, \boldsymbol{a}_n$ and $\boldsymbol{b}_1, \boldsymbol{b}_2, \dots, \boldsymbol{b}_n$ for the rows of A and of A^{-1}, respectively. There is a commutative diagram

$$\begin{tikzcd}
(\mathbb{C}^\times)^n \arrow[r, "\chi^{\boldsymbol{a}}"] \arrow[d, "\varphi_A"] & \mathbb{C}^\times \\
(\mathbb{C}^\times)^n \arrow[u, bend left, "\varphi_{A^{-1}}"] \arrow[ur, "\chi^{\boldsymbol{e}_1}"'] &
\end{tikzcd}$$

where φ_A and $\varphi_{A^{-1}}$ are the monomial isomorphisms given by $\boldsymbol{x} \mapsto \boldsymbol{x}^A = (\boldsymbol{x}^{\boldsymbol{a}_1}, \dots, \boldsymbol{x}^{\boldsymbol{a}_n})$ and $\boldsymbol{x} \mapsto \boldsymbol{x}^{A^{-1}} = (\boldsymbol{x}^{\boldsymbol{b}_1}, \dots, \boldsymbol{x}^{\boldsymbol{b}_n})$, respectively. Let $\boldsymbol{y} = (y_1, \dots, y_n)$ denote the coordinates of the torus below. For $\ell = 1, \dots, n$, set

$$f_\ell^A(\boldsymbol{y}) = \varphi_{A^{-1}}^* f_\ell = f_\ell(\boldsymbol{y}^{\boldsymbol{b}_1}, \dots, \boldsymbol{y}^{\boldsymbol{b}_n}) \in \mathbb{C}[\boldsymbol{y}^{\pm 1}],$$

so that $(\varphi_A)_* Z(\boldsymbol{f}) = Z(\boldsymbol{f}^A)$. Hence, $E_{\boldsymbol{a}}(\boldsymbol{f}) = E_{\boldsymbol{f}^A, \boldsymbol{e}_1}$ and so Lemma 3.7 implies that

$$\log \|E_{\boldsymbol{a}}(\boldsymbol{f})\|_{\sup} \le \sum_{j=1}^{n} \mathrm{MV}_{\boldsymbol{e}_1^\perp}(\{\pi_{\boldsymbol{e}_1}(\mathrm{N}(f_\ell^A))\}_{\ell \neq j}) \log \|f_j^A\|_{\sup}, \tag{3.10}$$

where $\mathrm{N}(f_\ell^A)$ is the Newton polytope of f_{ℓ^A}. We have $\pi_{\boldsymbol{e}_1}(\mathrm{N}(f_\ell^A)) = \widetilde{B}(\mathrm{N}(f_\ell)) \subset \widetilde{B}(Q_\ell)$ for the linear map $\widetilde{B}\colon \mathbb{R}^n \to \mathbb{R}^{n-1}$ given by $\widetilde{B}(\boldsymbol{x}) = (\langle \boldsymbol{x}, \boldsymbol{b}_2\rangle, \dots, \langle \boldsymbol{x}, \boldsymbol{b}_n\rangle)$. Let $\{\boldsymbol{v}_2, \dots, \boldsymbol{v}_n\}$ be an orthonormal basis of $\boldsymbol{a}^\perp$ and consider a second commutative diagram

$$\begin{tikzcd}
\mathbb{R}^n \arrow[r, "\widetilde{B}"] \arrow[d, "\pi_{\boldsymbol{a}}"'] & \mathbb{R}^{n-1} \arrow[d, "C"] \\
\boldsymbol{a}^\perp \arrow[r, "U"'] & \boldsymbol{a}^\perp
\end{tikzcd}$$

where C is the linear map defined by $C(y_2, \dots, y_n) = y_2\boldsymbol{v}_2 + \cdots + y_n\boldsymbol{v}_n$. It is easy to verify that $U \in \mathrm{GL}(\boldsymbol{a}^\perp)$ is uniquely determined by $\pi_{\boldsymbol{a}}$, $\widetilde{B}$ and C. Since C maps the canonical basis of $\mathbb{R}^{n-1}$ into an orthonormal basis of $\boldsymbol{a}^\perp$, it is an isometry between these two spaces. On the other hand, a straightforward computation shows that

$$U(\boldsymbol{v}_j) = \sum_{k=2}^{n} \langle \boldsymbol{v}_j, \boldsymbol{b}_k\rangle \boldsymbol{v}_k = \boldsymbol{b}_j, \qquad j = 2, \dots, n.$$

We note that $\boldsymbol{b}_2, \dots, \boldsymbol{b}_n$ is a basis of the $\mathbb{Z}$-module $\boldsymbol{a}^\perp \cap \mathbb{Z}^n$. The Brill-Gordan formula [Bou70, Chapitre 3, § 11, Proposition 15]) implies that $\mathrm{vol}(\boldsymbol{a}^\perp/(\boldsymbol{a}^\perp \cap \mathbb{Z}^n)) = \|\boldsymbol{a}\|_2$. Hence,

$$|\det(U)| = \mathrm{vol}(\boldsymbol{a}^\perp/(\boldsymbol{a}^\perp \cap \mathbb{Z}^n)) = \|\boldsymbol{a}\|_2.$$

Standard properties of the mixed volume imply that, for $j = 1, \dots, n$,

$$\begin{aligned}
\mathrm{MV}_{\boldsymbol{e}_1^\perp}(\{\widetilde{B}(Q_\ell)\}_{\ell\neq j}) &= \mathrm{MV}_{\boldsymbol{a}^\perp}(\{C \circ \widetilde{B}(Q_\ell)\}_{\ell \neq j}) \\
&= \mathrm{MV}_{\boldsymbol{a}^\perp}(\{U \circ \pi_{\boldsymbol{a}}(Q_\ell)\}_{\ell\neq j}) = \|\boldsymbol{a}\|_2 \, \mathrm{MV}_{\boldsymbol{a}^\perp}(\{\pi_{\boldsymbol{a}}(Q_\ell)\}_{\ell\neq j}).
\end{aligned}$$

In addition, φ_A gives an automorphism of $(S^1)^n$ and so $\|f_\ell^A\|_{\sup} = \|f_\ell\|_{\sup}$. We conclude that, when $\boldsymbol{a}$ is primitive,

$$\log \|E_{\boldsymbol{a}}(\boldsymbol{f})\|_{\sup} \le \|\boldsymbol{a}\|_2 \sum_{j=1}^{n} \mathrm{MV}_{\boldsymbol{a}^\perp}(\{\pi_{\boldsymbol{a}}(Q_\ell)\}_{\ell\neq j}) \log \|f_j\|_{\sup}. \tag{3.11}$$

Now let $\boldsymbol{a} \in \mathbb{Z}^n \setminus \{0\}$ be any vector. Choose a primitive $\boldsymbol{a}' \in \mathbb{Z}^n$ and $m \in \mathbb{Z}_{\ge 1}$ such that $\boldsymbol{a} = m\boldsymbol{a}'$. Using Proposition 3.3, we deduce that

$$E_{\boldsymbol{a}}(\boldsymbol{f})(z) = \pm \prod_{\omega \in \mu_m} E_{\boldsymbol{a}'}(\boldsymbol{f})(\omega z)$$

where μ_m denotes the set of m-th roots of 1. Hence $\|\boldsymbol{a}\|_2 = m\|\boldsymbol{a}'\|_2$, $\pi_{\boldsymbol{a}} = \pi_{\boldsymbol{a}'}$ and $\log \|E_{\boldsymbol{a}}(\boldsymbol{f})\|_{\sup} \le m \log \|E_{\boldsymbol{a}'}(\boldsymbol{f})\|_{\sup}$. The result follows from the bound (3.11) applied to $\boldsymbol{a}'$. □

Proof of Theorem 1.4. Let $\boldsymbol{a} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$. Applying Proposition 3.5 with $f_0 = z - \boldsymbol{x}^{\boldsymbol{a}}$ and $f_0 = z - \boldsymbol{x}^{-\boldsymbol{a}}$, we get that the product of the leading and the constant coefficients of $E_{\boldsymbol{a}}(\boldsymbol{f})$ is equal to

$$\pm \prod_{\boldsymbol{v}} \mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}}, \dots, \mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}}, \dots, f_n^{\boldsymbol{v}})^{|\langle \boldsymbol{v}, \boldsymbol{a}\rangle|}.$$

Recall also that $Z(E_{\boldsymbol{a}}(\boldsymbol{f})(z)) = \boldsymbol{\chi}_*^{\boldsymbol{a}}\big(Z(\boldsymbol{f})\big)$. Erdös-Turán's theorem (Theorem 1.1) then implies that

$$\Delta_{\mathrm{ang}}\big(\boldsymbol{\chi}_*^{\boldsymbol{a}}(Z(\boldsymbol{f})\big) \le c\sqrt{\frac{1}{D}\log\left(\frac{\|E_{\boldsymbol{a}}(\boldsymbol{f})(z)\|_{\sup}}{\prod_{\boldsymbol{v}} |\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})|^{\frac{|\langle \boldsymbol{v},\boldsymbol{a}\rangle|}{2}}}\right)},$$

with $c = 2.5619\dots$ Lemma 3.9 implies that $\log\left(\frac{\|E_{\boldsymbol{a}}(\boldsymbol{f})(z)\|_{\sup}}{\prod_{\boldsymbol{v}} |\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})|^{\frac{|\langle \boldsymbol{v},\boldsymbol{a}\rangle|}{2}}}\right)$ is bounded above by the quantity

$$\|\boldsymbol{a}\|_2 \log\left(\frac{\prod_{i=1}^n \|f_i\|_{\sup}^{D_{\boldsymbol{w},i}}}{\prod_{\boldsymbol{v}} |\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})|^{\frac{|\langle \boldsymbol{v},\boldsymbol{w}\rangle|}{2}}}\right),$$

for $\boldsymbol{w} = \frac{\boldsymbol{a}}{\|\boldsymbol{a}\|_2} \in S^{n-1}$. From the definitions of $\theta(Z(\boldsymbol{f}))$ and of the Erdös-Turán size $\eta(\boldsymbol{f})$ given in (2.2) and (1.3), respectively, we get

$$\theta(Z(\boldsymbol{f})) \le \min\{1, c\sqrt{\eta(\boldsymbol{f})}\}.$$

Applying Theorem 2.3 and the fact that the function $t^{\frac{2}{3}}(9 - \log(t))$ is monotonic increasing in the interval $(0,1]$, we deduce that

$$\begin{aligned}
\Delta_{\mathrm{ang}}(Z(\boldsymbol{f})) &\le 22n\left(\frac{8}{3}\right)^n \big(9 - \log(\min\{1, c\sqrt{\eta(\boldsymbol{f})}\})\big)^{\frac{2}{3}(n-1)} \min\{1, c\sqrt{\eta(\boldsymbol{f})}\}^{\frac{2}{3}} \\
&\le 22n\left(\frac{8}{3}\right)^n 2^{-\frac{2}{3}(n-1)}(18 + \log^+(\eta(\boldsymbol{f})^{-1}))^{\frac{2}{3}(n-1)} c^{\frac{2}{3}} \eta(\boldsymbol{f})^{\frac{1}{3}} \\
&\le 66\, n\, 2^n (18 + \log^+(\eta(\boldsymbol{f})^{-1}))^{\frac{2}{3}(n-1)} \eta(\boldsymbol{f})^{\frac{1}{3}},
\end{aligned}$$

which gives the bound for the angle discrepancy. For the radius discrepancy, we use the bounds given in Theorem 1.1, (2.2), and Theorem 2.3 to get, for $0 < \varepsilon < 1$,

$$\Delta_{\mathrm{rad}}(\boldsymbol{f}, \varepsilon) \le \sum_{j=1}^n \Delta_{\mathrm{rad}}(\boldsymbol{\chi}_*^{\boldsymbol{e}_j}(Z(\boldsymbol{f})), \varepsilon) \le \frac{2n}{\varepsilon}\,\eta(\boldsymbol{f}).$$

This concludes with the proof of the theorem. □

We next study a number of basic properties of the Erdös-Turán size. The following proposition shows that this notion generalizes the measure of polynomials that appears in the statement of Theorem 1.1.

Proposition 3.12. *Let* $d \ge 1$ *and* $f = a_0 + \dots + a_d x^d \in \mathbb{C}[x]$ *with* $a_0 a_d \neq 0$. *Then*

$$\eta(f) = \frac{1}{d}\log\left(\frac{\|f\|_{\sup}}{\sqrt{|a_0 a_d|}}\right).$$

Proof. The facet resultants of f are

$$\mathrm{Res}_{\boldsymbol{v}}(f^{\boldsymbol{v}}) \begin{cases} \pm a_0 & \text{for } \boldsymbol{v} = 1, \\ \pm a_d & \text{for } \boldsymbol{v} = -1. \end{cases}$$

Moreover, $D = \mathrm{MV}_{\mathbb{R}}([0,d]) = d$, $D_{\boldsymbol{w},1} = 1$ for $\boldsymbol{w} \in S^0 = \{\pm 1\}$, and $|\langle \boldsymbol{v}, \boldsymbol{w}\rangle| = 1$ for all $\boldsymbol{v}, \boldsymbol{w} \in \{\pm 1\}$. The formula for $\eta(f)$ then boils down to $\frac{1}{d}\log\left(\frac{\|f\|_{\sup}}{\sqrt{|a_0 a_d|}}\right)$. □

We denote by $\Delta^n = \mathrm{conv}(\boldsymbol{0}, \boldsymbol{e_1}, \dots, \boldsymbol{e_n})$ the standard simplex of $\mathbb{R}^n$.

Proposition 3.13. *Let* $\mathcal{A}_1,\dots,\mathcal{A}_n$ *be family of non-empty finite subsets of* $\mathbb{Z}^n$ *such that* $\mathrm{MV}(Q_1,\dots,Q_n)\geq 1$ *with* $Q_i=\mathrm{conv}(\mathcal{A}_i)$. *Let* $f_1,\dots,f_n\in\mathbb{C}[x_1^{\pm1},\dots,x_n^{\pm1}]$ *with* $\mathrm{supp}(f_i)\subset\mathcal{A}_i$ *and such that* $\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})\neq 0$ *for all* $\boldsymbol{v}\in\mathbb{Z}^n\setminus\{0\}$.

(1) $\eta(\boldsymbol{f})<\infty$.

(2) *Let* $\lambda_1,\dots,\lambda_n\in\mathbb{C}^\times$. *Then* $\eta(\lambda_1f_1,\dots,\lambda_nf_n)=\eta(f_1,\dots,f_n)$.

(3) *Let* $d_j\in\mathbb{Z}_{\geq1}$ *and* $\boldsymbol{b}_j\in\mathbb{Z}^n$ *such that* $\mathrm{N}(f_j)\subset d_j\Delta^n+\boldsymbol{b}_j$, $j=1,\dots,n$. *Then*

$$\eta(\boldsymbol{f})\leq\frac{1}{\mathrm{MV}(Q_1,\dots,Q_n)}\bigg((n+\sqrt{n})\Big(\prod_{j=1}^n d_j\Big)\sum_{j=1}^n\frac{\log\|f_j\|_{\sup}}{d_j}\\ +\sum_{\boldsymbol{v}}\frac{\|\boldsymbol{v}\|_2}{2}\log^+|\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})^{-1}|\bigg),$$

the second sum being taken over all primitive vectors $\boldsymbol{v}\in\mathbb{Z}^n$. *If* $f_1,\dots,f_n\in\mathbb{Z}[x_1^{\pm1},\dots,x_n^{\pm1}]$, *then*

$$\eta(\boldsymbol{f})\leq\frac{(n+\sqrt{n})\big(\prod_{j=1}^n d_j\big)}{\mathrm{MV}(Q_1,\dots,Q_n)}\sum_{j=1}^n\frac{\log\|f_j\|_{\sup}}{d_j}.$$

Proof. The statement of (1) is clear, since $\eta(\boldsymbol{f})$ is defined as the supremum of a continuous function over the compact set S^{n-1}.

For (2), let $\boldsymbol{a}\in\mathbb{Z}^n\setminus\{\boldsymbol{0}\}$. As explained in the proof of Theorem 1.4, the product of the leading and the constant coefficients of $E_{\boldsymbol{a}}(\boldsymbol{f})$ is equal to

$$\pm\prod_{\boldsymbol{v}}\mathrm{Res}_{\mathcal{A}_1^{\boldsymbol{v}},\dots,\mathcal{A}_n^{\boldsymbol{v}}}(f_1^{\boldsymbol{v}},\dots,f_n^{\boldsymbol{v}})^{|\langle\boldsymbol{v},\boldsymbol{a}\rangle|}.$$

Hence, the denominator in the definition of $\eta(\boldsymbol{f})$ is multihomogeneous in the coefficients of each f_j, of partial degrees equal to $\|\boldsymbol{a}\|_2{}^{-1}$ times those of $E_{\boldsymbol{a}}(\boldsymbol{f})$. Hence,

$$\frac{1}{\|\boldsymbol{a}\|_2}\deg_{f_j}(E_{\boldsymbol{a}}(\boldsymbol{f}))=\mathrm{MV}_{\boldsymbol{a}^\perp}\left(\pi_{\boldsymbol{a}}(Q_1),\dots,\pi_{\boldsymbol{w}}(Q_{j-1}),\pi_{\boldsymbol{a}}(Q_{j+1}),\dots,\pi_{\boldsymbol{a}}(Q_n)\right)=D_{\boldsymbol{w},j}$$

for $\boldsymbol{w}=\frac{\boldsymbol{a}}{\|\boldsymbol{a}\|_2}$, which implies the statement.

(3): let $\boldsymbol{w}\in S^{n-1}$. Then $\pi_{\boldsymbol{w}}(\mathrm{N}(f_j))\subset\pi_{\boldsymbol{w}}(d_j\Delta^n+\boldsymbol{b}_j)$. Due to the monotonicity of the mixed volume with respect to the inclusion, plus its properties of homogeneity and invariance under translation, we deduce that, for $j=1,\dots,n$,

$$\mathrm{MV}_{\boldsymbol{w}^\perp}(\{\pi_{\boldsymbol{w}}(\mathrm{N}(f_\ell))\}_{\ell\neq j})\leq\mathrm{MV}_{\boldsymbol{w}^\perp}(\{\pi_{\boldsymbol{w}}(d_\ell\Delta^n+\boldsymbol{b}_\ell)\}_{\ell\neq j})\\ \leq(n-1)!\Big(\prod_{\ell\neq j}d_\ell\Big)\mathrm{vol}_{\boldsymbol{w}^\perp}(\pi_{\boldsymbol{w}}(\Delta^n)).\tag{3.14}$$

The projected simplex $\pi_{\boldsymbol{w}}(\Delta^n)$ can be covered by the union of the projection of its facets. One the facets of Δ^n has $(n-1)$-dimensional volume equal to $\frac{\sqrt{n}}{(n-1)!}$, while the other n facets have $(n-1)$-dimensional volume equal to $\frac{1}{(n-1)!}$. Since the volume cannot increase under orthogonal projections, we have that

$$\mathrm{vol}_{\boldsymbol{w}^\perp}(\pi_{\boldsymbol{w}}(\Delta^n))\leq\frac{\sqrt{n}+n}{(n-1)!}.\tag{3.15}$$

In addition, $|\langle\boldsymbol{v},\boldsymbol{w}\rangle|\leq\|\boldsymbol{v}\|_2$ since $\boldsymbol{w}\in S^{n-1}$. Then, the first part of the statement follows from (3.14), (3.15) and the definition of $\eta(\boldsymbol{f})$. The second part follows from the fact that, if coefficients of the f_i's are integers, then the relevant facet resultants are nonzero integers and so their modulus are greater or equal to 1. ☐

Let us consider the statement of Proposition 3.13(3), in the classical dense case $Q_j = d_j\Delta^n$ for all j. In this situation, the only primitive vectors $\boldsymbol{v}$ to consider are $\boldsymbol{v} = \boldsymbol{e}_i$, $i = 1, \dots, n$, and $\boldsymbol{v} = \boldsymbol{e}_0 := -\sum_{i=1}^n \boldsymbol{e}_i$. Given $d_j \geq 1$, $j = 1, \dots, n$, we write $\mathrm{Res}_{d_1,\dots,d_n}$ for the resultant of n homogeneous polynomials in n variables of respective degrees $d_1, \dots, d_n$, as defined in [CLO05]. Given a system of polynomials $f_1, \dots, f_n \in \mathbb{C}[x_1, \dots, x_n]$ with $\deg(f_j) \leq d_j$ and $i = 0, \dots, n$, the facet polynomials $f_1^{\boldsymbol{e}_i}, \dots, f_n^{\boldsymbol{e}_i}$ form a system of n polynomials of degrees $d_1, \dots, d_n$. In particular, we can evaluate $\mathrm{Res}_{d_1,\dots,d_n}$ at these polynomials. If we assume that these resultants are nonzero, we obtain

$$\eta(\boldsymbol{f}) \leq (n+\sqrt{n}) \sum_{j=1}^n \frac{\log \|f_j\|_{\sup}}{d_j} + \frac{1}{2\prod_{j=1}^n d_j}\Big(\sqrt{n}\log^+|\mathrm{Res}_{d_1,\dots,d_n}(f_1^{\boldsymbol{e}_0}, \dots, f_n^{\boldsymbol{e}_0})^{-1}| + \sum_{i=1}^n \log^+|\mathrm{Res}_{d_1,\dots,d_n}(f_1^{\boldsymbol{e}_i}, \dots, f_n^{\boldsymbol{e}_i})^{-1}|\Big).$$

In particular, if $f_1, \dots, f_n \in \mathbb{Z}[x_1, \dots, x_n]$, then

$$\eta(\boldsymbol{f}) \leq (n+\sqrt{n}) \sum_{j=1}^n \frac{\log \|f_j\|_{\sup}}{d_j}.$$

4. Asymptotic equidistribution

We will apply here the results in the previous sections to study the asymptotic distribution of the roots of a sequence of systems of Laurent polynomials over $\mathbb{Z}$ and of random systems of Laurent polynomials over $\mathbb{C}$.

First, we will consider polynomials over $\mathbb{Z}$. Let $Q_i, \dots, Q_n \subset \mathbb{R}^n$ be a family of lattice polytopes such that $\mathrm{MV}(Q_1, \dots, Q_n) \geq 1$. For each $d \geq 1$ and $i = 1, \dots, n$, consider the finite subset of $\mathbb{Z}^n$ given by

$$\mathcal{A}_{i,d} = dQ_i \cap \mathbb{Z}^n. \tag{4.1}$$

Proposition 4.2. *For $d \geq 1$ let $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ be a family of Laurent polynomials in $\mathbb{Z}[x_1^{\pm1}, \dots, x_n^{\pm1}]$ such that $\mathrm{supp}(f_{i,d}) \subset dQ_i$ and $\mathrm{Res}_{\mathcal{A}_{1,d}^{\boldsymbol{v}},\dots,\mathcal{A}_{n,d}^{\boldsymbol{v}}}(f_{1,d}^{\boldsymbol{v}}, \dots, f_{n,d}^{\boldsymbol{v}}) \neq 0$ for all $\boldsymbol{v} \in \mathbb{Z}^n \setminus \{\boldsymbol{0}\}$. Then there is a constant $c_1 > 0$ which does not depend on d such that*

$$\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) \leq c_1 \left(\frac{\sum_{i=1}^n \log\|f_{i,d}\|_{\sup}}{d}\right)^{\frac{1}{3}} \left(1 + \log^+\left(\frac{d}{\sum_{i=1}^n \log\|f_{i,d}\|_{\sup}}\right)\right)^{\frac{2}{3}(n-1)}$$

and, for any $0 < \varepsilon < 1$,

$$\Delta_{\mathrm{rad}}(Z(\boldsymbol{f}_d), \varepsilon) \leq c_1 \frac{\sum_{i=1}^n \log\|f_{i,d}\|_{\sup}}{\varepsilon d}.$$

Proof. This follows easily from Theorem 1.4 and Proposition 3.13(3). ☐

Proof of Theorem 1.7. Let notations be as in the statement of Theorem 1.7. In particular, $\nu_{Z(\boldsymbol{f}_d)}$ is the discrete measure associated to $Z(\boldsymbol{f}_d)$, and ν is the measure on $(\mathbb{C}^\times)^n$ induced by the Haar probability measure on $(S^1)^n$.

We have to show that, for every continuous function with compact support h,

$$\lim_{d\to\infty}\int_{(\mathbb{C}^\times)^n} h\,\mathrm{d}\nu_{Z(\boldsymbol{f}_d)} = \int_{(\mathbb{C}^\times)^n} h\,\mathrm{d}\nu.$$

It is enough to prove the statement for the characteristic function h_U of the open sets of the form

$$U = \{(z_1,\dots,z_n)\in(\mathbb{C}^\times)^n \mid r_{1,j} < |z_j| < r_{2,j},\ \alpha_j < \arg(z_j) < \beta_j \text{ for all } j\}, \tag{4.3}$$

with $0\le r_{1,j} < r_{2,j}\le\infty$, $r_{i,j}\ne 1$ and $-\pi<\alpha_j<\beta_j\le\pi$, since any continuous function with compact support can be uniformly approximated by a linear combinations of these.

Consider first the case when $U\cap(S^1)^n=\emptyset$. Due to the conditions imposed on the numbers $r_{i,j}$, there exists $\varepsilon>0$ such that U is disjoint with the set $\{\boldsymbol{\xi}\in\mathbb{C}^n \mid 1-\varepsilon < |\xi_j| < (1-\varepsilon)^{-1} \text{ for all } j\}$. Hence,

$$\int_{(\mathbb{C}^\times)^n} h_U\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} = \frac{\deg(Z(\boldsymbol{f}_d)|_U)}{d^n\,\mathrm{MV}(\boldsymbol{Q})} \le \Delta_{\mathrm{rad}}(\boldsymbol{f}_d,\varepsilon),$$

where we set $\boldsymbol{Q}=(Q_1,\dots,Q_n)$ and $Z(\boldsymbol{f}_d)|_U=\sum_{\boldsymbol{\xi}\in|Z(\boldsymbol{f}_d)|\cap U} m_{\boldsymbol{\xi}}\boldsymbol{\xi}$. Proposition 4.2 implies that this integral goes to 0 for $d\to\infty$, which proves the statement in this case, since $\int_{\mathbb{C}^n} h_U\,\mathrm{d}\nu=0$.

Consider now the case when $U\cap(S^1)^n\ne\emptyset$. Set $\overline{U}=\{\boldsymbol{z}\mid \alpha_j<\arg(z_j)<\beta_j \text{ for all } j\}$. Then

$$\begin{aligned}\int_{(\mathbb{C}^\times)^n} h_U\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} - \int_{(\mathbb{C}^\times)^n} h_U\,\mathrm{d}\nu &= \int_{(\mathbb{C}^\times)^n} h_U\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} - \prod_{j=1}^n\frac{\beta_j-\alpha_j}{2\pi}\\ &= \int_{(\mathbb{C}^\times)^n}\Big(h_{\overline{U}} - \prod_{j=1}^n\frac{\beta_j-\alpha_j}{2\pi}\Big)\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} - \int_{(\mathbb{C}^\times)^n} h_{\overline{U}\setminus U}\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)}.\end{aligned}$$

We have

$$\int_{(\mathbb{C}^\times)^n}\Big|h_{\overline{U}} - \prod_{j=1}^n\frac{\beta_j-\alpha_j}{2\pi}\Big|\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} \le \Big|\frac{\deg(Z(\boldsymbol{f}_d)_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{d^n\,\mathrm{MV}(\boldsymbol{Q})} - \prod_{j=1}^n\frac{\beta_j-\alpha_j}{2\pi}\Big| \le \Delta_{\mathrm{ang}}(\boldsymbol{f}_d).$$

Again, Proposition 4.2 implies that this integral goes to 0 for $d\to\infty$. On the other hand, $\overline{U}\setminus U$ is a union of a finite number of subsets U_l of the form (4.3) such that $U_l\cap(S^1)^n=\emptyset$ for all l. By the previous considerations, $\int_{(\mathbb{C}^\times)^n} h_{U_l}\mathrm{d}\delta_{Z(\boldsymbol{f}_d)}\to_d 0$ and so $\int_{(\mathbb{C}^\times)^n} h_{\overline{U}\setminus U}\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)}\to_d 0$. Hence

$$\lim_{d\to\infty}\int_{(\mathbb{C}^\times)^n} h_U\,\mathrm{d}\delta_{Z(\boldsymbol{f}_d)} = \prod_{j=1}^n\frac{\beta_j-\alpha_j}{2\pi} = \int_{\mathbb{C}^n} h_U\,\mathrm{d}\nu = 0,$$

which concludes the proof. □

We will now consider random systems of Laurent polynomials with complex coefficients. To explain and prove our results, we have to consider metrics and measures on projective spaces over $\mathbb{C}$. Let $N\ge1$ and consider the standard Riemannian structure on $\mathbb{C}^N$ induced by the Euclidean norm $\|\cdot\|_2$. Let $S^{2N-1}=\{\boldsymbol{z}\in\mathbb{C}^N \mid \|\boldsymbol{z}\|_2=1\}$ be the unit sphere with the induced Riemannian structure. The map $S^{2N-1}\to\mathbb{P}(\mathbb{C}^N)$ given by $(z_0,\dots,z_{N-1})\mapsto(z_0:\cdots:z_{N-1})$ gives a principal bundle with fiber S^1. The

Fubini-Study metric on $\mathbb{P}(\mathbb{C}^N)$ is defined as the unique Riemannian structure such that this map is a Riemannian submersion, see [KN69] for details.

The geodesics of $\mathbb{P}(\mathbb{C}^N)$ coincide with the lines. Hence, we can define a distance between two points $\boldsymbol{z}_1$ and $\boldsymbol{z}_2$ as the length of the line segment joining them, and we will denote it by $\operatorname{dist}_{\mathrm{FS}}(\boldsymbol{z}_1, \boldsymbol{z}_2)$. However, it will be more convenient to consider the distance function $\operatorname{dist} := \sin(\operatorname{dist}_{\mathrm{FS}})$. This function can be computed with the formula

$$\operatorname{dist}(\boldsymbol{z}_1, \boldsymbol{z}_2) = \sqrt{1 - \Big(\frac{|\langle \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2\rangle|}{\|\widetilde{\boldsymbol{z}}_1\|_2 \|\widetilde{\boldsymbol{z}}_2\|_2}\Big)^2} \tag{4.4}$$

for any choice of representatives $\widetilde{\boldsymbol{z}}_i \in \mathbb{C}^N \setminus \{\boldsymbol{0}\}$, $i = 1, 2$.

Lemma 4.5. *Let* $\boldsymbol{z}_1, \boldsymbol{z}_2 \in \mathbb{P}(\mathbb{C}^N)$ *with* $\widetilde{\boldsymbol{z}}_i \in \mathbb{C}^N$, $i = 1, 2$, *representatives of these points, and such that* $\|\widetilde{\boldsymbol{z}}_2\|_2 = 1$. *Then* $\operatorname{dist}(\boldsymbol{z}_1, \boldsymbol{z}_2) \le \|\widetilde{\boldsymbol{z}}_2 - \widetilde{\boldsymbol{z}}_1\|_2$.

Proof. We have that

$$\|\widetilde{\boldsymbol{z}}_2 - \widetilde{\boldsymbol{z}}_1\|_2^2 = \langle \widetilde{\boldsymbol{z}}_2 - \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2 - \widetilde{\boldsymbol{z}}_1\rangle = 1 + \|\widetilde{\boldsymbol{z}}_1\|_2^2 - 2\operatorname{Re}(\langle \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2\rangle) \ge 1 + \|\widetilde{\boldsymbol{z}}_1\|_2^2 - 2|\langle \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2\rangle|.$$

On the other hand, the formula (4.4) gives $\operatorname{dist}(\boldsymbol{z}_1, \boldsymbol{z}_2)^2 = 1 - \big(\frac{|\langle \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2\rangle|}{\|\widetilde{\boldsymbol{z}}_1\|_2}\big)^2$. Hence,

$$\|\widetilde{\boldsymbol{z}}_2 - \widetilde{\boldsymbol{z}}_1\|_2^2 - \operatorname{dist}(\boldsymbol{z}_1, \boldsymbol{z}_2)^2 = \Big(\|\widetilde{\boldsymbol{z}}_1\|_2 - \frac{|\langle \widetilde{\boldsymbol{z}}_1, \widetilde{\boldsymbol{z}}_2\rangle|}{\|\widetilde{\boldsymbol{z}}_1\|_2}\Big)^2 \ge 0,$$

which proves the statement. □

We will need the following Łojasiewicz inequality for a hypersurface of a complex projective space. For a homogeneous polynomial $f \in \mathbb{C}[\boldsymbol{x}]$ of degree d, and a point $\boldsymbol{z} \in \mathbb{P}(\mathbb{C}^N)$, the value

$$\frac{|f(\boldsymbol{z})|}{\|\boldsymbol{z}\|_2^d}$$

is well-defined. For a subset $E \subset \mathbb{P}(\mathbb{C}^N)$, we write $\operatorname{dist}(\boldsymbol{z}, E)$ for the distance between $\boldsymbol{z}$ and E.

Lemma 4.6. *Let* $f \in \mathbb{C}[x_0, \dots, x_{N-1}]$ *be a homogeneous polynomial of degree* $d \ge 0$ *and* $\boldsymbol{z} \in \mathbb{P}(\mathbb{C}^N)$. *Then*

$$\frac{|f(\boldsymbol{z})|}{\|\boldsymbol{z}\|_2^d} \ge \Big(\sup_{\boldsymbol{x} \in \mathbb{P}(\mathbb{C}^N)} \frac{|f(\boldsymbol{x})|}{\|\boldsymbol{x}\|_2^d}\Big) \operatorname{dist}(\boldsymbol{z}, V(f))^d.$$

Proof. Let $\boldsymbol{z}, \boldsymbol{x} \in \mathbb{P}(\mathbb{C}^N)$ such that $\boldsymbol{x} \notin V(f)$. Let $\widetilde{\boldsymbol{z}}, \widetilde{\boldsymbol{x}}$ be representatives of these points in the sphere S^{2N-1} and set $f_{\widetilde{\boldsymbol{x}}}(t) = f(\widetilde{\boldsymbol{z}} + t\widetilde{\boldsymbol{x}}) \in \mathbb{C}[t]$. This is a univariate polynomial of degree d with leading coefficient $f(\widetilde{\boldsymbol{x}})$. Then, there exist $\xi_j \in \mathbb{C}$, $j = 1, \dots, d$, such that $f_{\widetilde{\boldsymbol{x}}} = f(\widetilde{\boldsymbol{x}}) \prod_j (t - \xi_j)$ and so

$$|f(\widetilde{\boldsymbol{z}})| = |f_{\widetilde{\boldsymbol{x}}}(0)| = |f(\widetilde{\boldsymbol{x}})| \prod_j |\xi_j|.$$

For each j, we have that $\widetilde{\boldsymbol{z}} + \xi_j \widetilde{\boldsymbol{x}} \in V(f)$. Using Lemma 4.5, we deduce that

$$|\xi_j| = \|(\widetilde{\boldsymbol{z}} + \xi_j \widetilde{\boldsymbol{x}}) - \widetilde{\boldsymbol{z}}\|_2 \ge \operatorname{dist}(\boldsymbol{z}, \widetilde{\boldsymbol{z}} + \xi_j \widetilde{\boldsymbol{x}}) \ge \operatorname{dist}(\boldsymbol{z}, V(f)).$$

We deduce that

$$\frac{|f(\boldsymbol{z})|}{\|\boldsymbol{z}\|_2^d} = |f(\widetilde{\boldsymbol{z}})| \ge |f(\widetilde{\boldsymbol{x}})| \operatorname{dist}(\boldsymbol{z}, V(f))^d = \frac{|f(\boldsymbol{x})|}{\|\boldsymbol{x}\|_2^d} \operatorname{dist}(\boldsymbol{z}, V(f))^d.$$

Since this holds for all $\boldsymbol{x} \notin V(f)$, the result follows. □

Let μ_{FS} denote the measure on $\mathbb{P}(\mathbb{C}^N)$ induced by the Fubini-Study metric. Then $\mu_{\mathrm{FS}}(\mathbb{P}(\mathbb{C}^N)) = \frac{\pi^{N-1}}{(N-1)!}$. We will consider the normalized measure given by

$$\mu = \frac{(N-1)!}{\pi^{N-1}}\mu_{\mathrm{FS}}.$$

A result of Beltrán and Pardo [BP07, Theorem 1] shows that, for a hypersurface $H \subset \mathbb{P}(\mathbb{C}^N)$ of degree d, the normalized measure of the tube around H of radius $\rho \geq 0$ is bounded above by

$$15d(N-1)^2\rho^2.$$

Applying this result, we deduce the following bound for the volume of the set where a polynomial can take small values. For $\delta > 0$ and a homogeneous polynomial $f \in \mathbb{C}[\boldsymbol{x}]$, we consider the subset of $\mathbb{P}(\mathbb{C}^N)$ given by

$$V(f)_\delta = \Big\{ \boldsymbol{z} \in \mathbb{P}(\mathbb{C}^N) \Big| \frac{|f(\boldsymbol{z})|}{\|\boldsymbol{z}\|_2^d} < \delta \Big\}. \tag{4.7}$$

Proposition 4.8. *Let $\delta > 0$ and $f \in \mathbb{C}[x_0, \dots, x_{N-1}]$ a homogeneous polynomial of degree $d \geq 1$. Then*

$$\mu(V(f)_\delta) \leq 15dN^3\Big(\frac{\delta}{\|f\|_{\sup}}\Big)^{\frac{2}{d}}.$$

In particular, if $f \in \mathbb{Z}[x_0, \dots, x_{N-1}]$, then $\mu(V(f)_\delta) \leq 15dN^3\delta^{\frac{2}{d}}$.

Proof. Let $\boldsymbol{z} \in V(f)_\delta$. Using Lemma 4.6, we deduce that

$$\delta > \bigg(\sup_{\boldsymbol{x}} \frac{|f(\boldsymbol{x})|}{\|\boldsymbol{x}\|_2^d}\bigg) \operatorname{dist}(\boldsymbol{z}, V(f))^d \geq \frac{\|f\|_{\sup}}{N^{\frac{d}{2}}} \operatorname{dist}(\boldsymbol{z}, V(f))^d.$$

Hence,

$$\operatorname{dist}(\boldsymbol{z}, V(f)) < N^{\frac{1}{2}}\Big(\frac{\delta}{\|f\|_{\sup}}\Big)^{\frac{1}{d}}$$

and so $V(f)_\delta$ is contained in the tube around $V(f)$ of radius $N^{\frac{1}{2}}\big(\frac{\delta}{\|f\|_{\sup}}\big)^{\frac{1}{d}}$. The first part of the result follows then from the Beltrán–Pardo bound for the volume of this tube. The second part follows from the fact that $\|f\|_{\sup} \geq |f| \geq 1$, because of the inequality (3.6) and the fact that the coefficients of f are integer numbers. □

Let us keep notations as above and set $\boldsymbol{\mathcal{A}}_d = (\mathcal{A}_{1,d}, \dots, \mathcal{A}_{n,d})$ with $\mathcal{A}_{i,d}$ as in (4.1). Each point of the projective space $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ can be identified with a system $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ of Laurent polynomials such that $\operatorname{supp}(f_{i,d}) \subset dQ_i$, $i = 1, \dots, n$, modulo a multiplicative scalar. The associated cycle $Z(\boldsymbol{f}_d)$ is well-defined, since it does not depend on this multiplicative scalar.

Set μ_d for the normalized Fubini-Study measure on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ and let $g_d \colon \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}) \to \mathbb{R}_{\geq 0}$ be a probability density function, that is, a μ_d-measurable function with

$$\int_{\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})} g_d \, \mathrm{d}\mu_d = 1.$$

Let $\boldsymbol{f}_d$ be a random system of Laurent polynomials with $\operatorname{supp}(f_{i,d}) \subset dQ_i$, $i = 1, \dots, n$, distributed according to the probability law given by g_d with respect to μ_d. We can then consider the angle discrepancy of $Z(\boldsymbol{f}_d)$ and, for $0 < \varepsilon < 1$, the radius discrepancy of $Z(\boldsymbol{f}_d)$ with respect to ε, as random variables on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$. We denote by $\mathbb{E}(\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)); g_d)$ and $\mathbb{E}(\Delta_{\mathrm{rad}}(Z(\boldsymbol{f}_d), \varepsilon); g_d)$ the expected value of these random variables.

Theorem 4.9. *For each* $d \geq 1$*, let* g_d *be a probability density function on* $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ *and* $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ *a random system of Laurent polynomials with* $\mathrm{supp}(f_{i,d}) \subset dQ_i$, $i = 1, \dots, n$*, distributed according to the probability law given by* g_d *with respect to* μ_d*. Assume that the sequence* $(g_d)_{d \geq 1}$ *is uniformly bounded. Then there is a constant* $c_2 > 0$ *which does not depend on* d *such that*

$$\mathbb{E}(\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)); g_d) \leq c_2 \, \frac{\log(d+1)^{\frac{2}{3}n - \frac{1}{3}}}{d^{\frac{1}{3}}}$$

and, for any $0 < \varepsilon < 1$,

$$\mathbb{E}(\Delta_{\mathrm{rad}}(Z(\boldsymbol{f}_d), \varepsilon); g_d) \leq c_2 \, \frac{\log(d+1)}{\varepsilon d}.$$

In particular, $\lim_{d \to \infty} \mathbb{E}(\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)); g_d) = 0$ *and* $\lim_{d \to \infty} \mathbb{E}(\Delta_{\mathrm{rad}}(Z(\boldsymbol{f}_d), \varepsilon); g_d) = 0$.

Proof. We first estimate the expected value of the angular discrepancy, which is given by the formula

$$\mathbb{E}(\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)); g_d) = \int_{\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})} \Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) g_d(\boldsymbol{f}) \, \mathrm{d}\mu_d.$$

Consider the Minkowski sum $Q = \sum_{i=1}^n Q_i$, which is a lattice polytope on $\mathbb{R}^n$ of dimension n because of the assumption that the mixed volume of $Q_1, \dots, Q_n$ is positive. For each primitive vector $\boldsymbol{v} \in \mathbb{Z}^n$ which is an inner normal to a facet of Q, consider the facet resultant

$$R_{\boldsymbol{v},d} = \mathrm{Res}_{\mathcal{A}_{1,d}^{\boldsymbol{v}}, \dots, \mathcal{A}_{n,d}^{\boldsymbol{v}}} \in \mathbb{Z}[\boldsymbol{u}_1, \dots, \boldsymbol{u}_n],$$

where $\boldsymbol{u}_i$ is a group of $\#\mathcal{A}_{i,d}$ variables. Proposition 3.13(2) implies that its total degree is bounded by $\deg(R_{\boldsymbol{v},d}) = c_{\boldsymbol{v}} d^{n-1}$ for a constant $c_{\boldsymbol{v}}$ independent of d. Its total number of variables is $\#\boldsymbol{\mathcal{A}}_d = \sum_i \mathcal{A}_{i,d} = \sum_i dQ_i \cap \mathbb{Z}^n$. This number can be bounded by $c_3 d^n$ for a constant c_3 independent of d.

Set $\delta_{\boldsymbol{v},d} = d^{-2n \deg(R_{\boldsymbol{v},d})}$. Consider the subset $V(R_{\boldsymbol{v},d})_{\delta_{\boldsymbol{v},d}} \subset \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ as defined in (4.7) and put

$$U_d = \bigcup_{\boldsymbol{v}} V(R_{\boldsymbol{v},d})_{\delta_{\boldsymbol{v},d}},$$

the union being over all primitive inner normal vectors to facets of Q. Using the fact that $0 \leq \Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) \leq 1$ and the hypothesis that the functions g_d are uniformly bounded, we deduce that

$$0 \leq \int_{U_d} \Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) g_d(\boldsymbol{f}) \, \mathrm{d}\mu_d \leq \Big(\sup_{\boldsymbol{f}_d} g_d(\boldsymbol{f}_d) \Big) \mu_d(U_d) \leq c_4 \mu_d(U_d)$$

for a constant c_4 independent of d. By Proposition 4.8,

$$\begin{aligned} \mu_d(U_d) \leq \sum_{\boldsymbol{v}} \mu_d(V(R_{\boldsymbol{v},d})_{\delta_{\boldsymbol{v},d}}) &\leq \sum_{\boldsymbol{v}} 15 \deg(R_{\boldsymbol{v},d}) \big(\#\boldsymbol{\mathcal{A}}_d\big)^3 \delta_{\boldsymbol{v},d}^{\frac{2}{\deg(R_{\boldsymbol{v},d})}} \\ &\leq 15 \Big(\sum_{\boldsymbol{v}} c_{\boldsymbol{v}} d^{n-1} \Big) (c_3 d^n)^3 d^{-4n} = c_5 d^{-1}, \end{aligned} \tag{4.10}$$

with $c_5 = 15 \big(\sum_{\boldsymbol{v}} c_{\boldsymbol{v}} \big) c_3^3$. Hence, $\mu_d(U_d) \to_d 0$ and so $\int_{U_d} \Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) g_d(\boldsymbol{f}_d) \, \mathrm{d}\mu_d \to_d 0$ as well.

Let $\boldsymbol{f}_d \in \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}) \setminus U_d$ and choose a representative $\widetilde{\boldsymbol{f}}_d = (\widetilde{f}_{1,d}, \dots, \widetilde{f}_{n,d}) \in \mathbb{C}^{\boldsymbol{\mathcal{A}}_d} \setminus \{\boldsymbol{0}\}$ with $\|\widetilde{\boldsymbol{f}}_d\|_2 = 1$. By Proposition 3.13(2), $\eta(\boldsymbol{f}_d) = \eta(\widetilde{\boldsymbol{f}}_d)$. Note that the Minkowski sum $\sum_i dQ_i$ coincides with dQ. Hence, the only non trivial facet resultants of the family of finite sets $\mathcal{A}_{1,d}, \dots, \mathcal{A}_{n,d}$ are those of the form $R_{\boldsymbol{v},d}$ as considered above. As before, denote with Δ^n the standard simplex of $\mathbb{R}^n$. Choose $e \geq 1$ and $\boldsymbol{b}_i \in \mathbb{Z}^n$ such that $Q_i \subset e\Delta^n + \boldsymbol{b}_i$ for all i. Hence, $d\mathcal{A}_{i,d} \subset deQ_i + d\boldsymbol{b}_i$ for all i. Proposition 3.13(3) implies that there is a constant c_6 independent of d such that

$$\begin{aligned}\eta(\boldsymbol{f}_d) &\leq \frac{1}{d^n \operatorname{MV}(\boldsymbol{Q})} \Big((de)^{n-1}(n+\sqrt{n}) \sum_{i=1}^{n} \log \|\widetilde{f}_{i,d}\|_{\sup} + \frac{1}{2} \sum_{\boldsymbol{v}} \|\boldsymbol{v}\|_2 \log^+ |R_{\boldsymbol{v},d}(\boldsymbol{f}_d^{\boldsymbol{v}})^{-1}| \Big) \\ &\leq \frac{1}{d^n \operatorname{MV}(\boldsymbol{Q})} \Big((de)^{n-1}(n+\sqrt{n}) \sum_{i=1}^{n} \log(\#\mathcal{A}_{i,d}) + n \sum_{\boldsymbol{v}} \|\boldsymbol{v}\|_2 \deg(R_{\boldsymbol{v},d}) \log(d) \Big) \\ &\leq c_6 \frac{\log(d+1)}{d},\end{aligned}$$

the second and fourth sums being over the primitive inner normals $\boldsymbol{v}$ to the facets of Q. Here, we used the fact that $\|\widetilde{f}_{i,d}\|_{\sup} \leq \#\mathcal{A}_{i,d}$ for $\widetilde{\boldsymbol{f}}_d$ in the unit sphere of $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$, the definition of the set U_d, the bound $\#\mathcal{A}_{i,d} \leq \#\boldsymbol{\mathcal{A}}_d \leq c_3 d^n$ and the inequality $\deg(R_{\boldsymbol{v},d}) \leq c_{\boldsymbol{v}} d^{n-1}$ that we explained before.

Using Theorem 1.4 and the fact that the function $t^{\frac{1}{3}} \log\left(\frac{\alpha}{t}\right)^{\frac{n-1}{3}}$ is increasing for small values of $t > 0$, we deduce that, for $\boldsymbol{f}_d \in \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}) \setminus U_d$,

$$\begin{aligned}\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) &\leq c_7 \eta(\boldsymbol{f}_d)^{\frac{1}{3}} \log\left(\frac{c_8}{\eta(\boldsymbol{f}_d)}\right)^{\frac{2}{3}(n-1)} \\ &\leq c_9 \Big(\frac{\log(d+1)}{d}\Big)^{\frac{1}{3}} \log\left(\frac{d}{\log(d+1)}\right)^{\frac{2}{3}(n-1)} \leq c_{10} \, \frac{\log(d+1)^{\frac{2}{3}n-\frac{1}{3}}}{d^{\frac{1}{3}}}\end{aligned}$$

for suitable constants c_7, c_8, c_9 and c_{10}. This proves the first part of the statement. For the radius discrepancy, we proceed in a similar way: given $\varepsilon > 0$, we write $\mathbb{E}(\Delta_{\mathrm{rad}}(\boldsymbol{f}_d, \varepsilon); g_d)$ as an integral, which we split into two parts. We bound the first using that $0 \leq \Delta_{\mathrm{rad}}(\boldsymbol{f}_d, \varepsilon) \leq 1$ and the estimate (4.10), while the second integral can be bounded by applying Theorem 1.4. □

Proof of Theorem 1.8. The proof is similar to the one given for Theorem 1.7. Write $\nu_d = \frac{\mathbb{E}(Z(\boldsymbol{f}_d); g_d)}{d^n \operatorname{MV}(\boldsymbol{Q})}$ for short, where $\mathbb{E}(Z(\boldsymbol{f}_d); g_d)$ is the expected zero density measure of $\boldsymbol{f}_d$. To prove the statement, it is enough to show that, for all subsets U as in (4.3),

$$\lim_{d \to \infty} \nu_d(U) = \nu(U \cap (S^1)^n) = \begin{cases} \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi} & \text{if } U \cap (S^1)^n \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

If $U \cap (S^1)^n = \emptyset$, then there exists $\varepsilon > 0$ such that

$$\deg(Z(\boldsymbol{f}_d)|_U) \leq \deg(Z(\boldsymbol{f}_d)) \Delta_{\mathrm{rad}}(\boldsymbol{f}_d, \varepsilon) \leq d^n \operatorname{MV}(\boldsymbol{Q}) \Delta_{\mathrm{rad}}(\boldsymbol{f}_d, \varepsilon)$$

and so $\nu_d(U) \leq \mathbb{E}(\Delta_{\mathrm{rad}}(\boldsymbol{f}_d, \varepsilon); g_d)$. Theorem 4.9 then implies that $\lim_{d\to\infty} \nu_d(U) = 0 = \nu(U)$.

If $U \cap (S^1)^n \neq \emptyset$, we set $\overline{U} = \{\boldsymbol{z} \mid \alpha_j < \arg(z_j) < \beta_j \text{ for all } j\}$, and then we have

$$\nu_d(U) - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi} = \Big(\nu_d(\overline{U}) - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi}\Big) - \nu_d(\overline{U} \setminus U).$$

Set $R_d = \prod_{\boldsymbol{v}} R_{\boldsymbol{v},d}$ for the product of all non-trivial facet resultants of the family of finite sets $\mathcal{A}_1, \dots, \mathcal{A}_n$. Then

$$\Big|\nu_d(\overline{U}) - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi}\Big| = \int_{\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})\setminus V(R_d)} \Big|\frac{\deg(Z(\boldsymbol{f}_d)_{\boldsymbol{\alpha},\boldsymbol{\beta}})}{d^n \operatorname{MV}(\boldsymbol{Q})} - \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi}\Big| g_d(\boldsymbol{f}_d)\, \mathrm{d}\mu_d$$
$$\leq \int_{\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})} \Delta_{\mathrm{ang}}(\boldsymbol{f}_d) g_d(\boldsymbol{f}_d)\, \mathrm{d}\mu_d.$$

We have that $\overline{U} \setminus U$ is a union of a finite number of subsets U_l of the form (4.3) such that $U_l \cap (S^1)^n = \emptyset$ for all l. By the previous considerations, $\lim_{d\to\infty} \nu_d(U_l) = 0$ and so $\lim_{d\to\infty} \nu_d(\overline{U} \setminus U) = 0$. Theorem 4.9 then implies that

$$\lim_{d\to\infty} \nu_d(U) = \lim_{d\to\infty} \nu_d(\overline{U}) = \prod_{j=1}^{n} \frac{\beta_j - \alpha_j}{2\pi} = \nu(U).$$

□

Remark 4.11. It is not clear to us whether the upper bound in Proposition 4.8 for the volume of the set $V(f)_\delta$ is sharp or not. It would be interesting to clarify this point, as a qualititive improvement on this bound might enlarge the range of applicability of theorems 4.9 and 1.8.

Remark 4.12. In some situations, it might be interesting to consider probability distributions on the complex linear space $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$ rather than on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$. For a point $\boldsymbol{f}_d \in \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$, the associated cycle $Z(\boldsymbol{f}_d)$ does not depend on the choice of a representative in $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$ for this point and, *a fortiori*, the same holds for the angle and radius discrepancies of $Z(\boldsymbol{f}_d)$. Hence, one might consider random variables on $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$ arising from this cycle as random variables on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$, by applying Federer's coarea formula (see for instance [BP07, Theorem 20]).

In precise terms, the normal Jacobian of the map $\varpi\colon \mathbb{C}^{\boldsymbol{\mathcal{A}}_d} \setminus \{\boldsymbol{0}\} \to \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ with respect to the Euclidean structure on $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d} \setminus \{\boldsymbol{0}\}$ and the Fubini-Study one on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ is given, for $\boldsymbol{g} \in \mathbb{C}^{\boldsymbol{\mathcal{A}}_d} \setminus \{\boldsymbol{0}\}$, by

$$\mathrm{NJ}_{\boldsymbol{g}}\, \varpi = \|\boldsymbol{g}\|^{-2N_d}.$$

with $N_d = \#\boldsymbol{\mathcal{A}}_d - 1$. Given a probability density function $G_d\colon \mathbb{C}^{\boldsymbol{\mathcal{A}}} \to \mathbb{R}$, one might derive a corresponding probability density function on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d})$ by integrating along the fibers of ϖ as

$$g_d(\boldsymbol{f}_d) = \frac{\pi^{N_d}}{N_d!} \int_{\varpi^{-1}(\boldsymbol{f}_d)} G_d(\boldsymbol{g}) \|\boldsymbol{g}\|_2^{2N_d}\, \mathrm{d}\varpi^{-1}(\boldsymbol{f}_d), \tag{4.13}$$

where $\mathrm{d}\varpi^{-1}(\boldsymbol{f}_d)$ is the volume form of the fiber $\varpi^{-1}(\boldsymbol{f}_d)$. The probability distribution given by G_d of, for instance, the angle discrepancy, can then be computed, for any Borel subset $I \subset [0,1]$, as

$$\operatorname{Prob}(\Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) \in I; G_d) = \int_{\Delta_{\mathrm{ang}}^{-1}(I)} g_d(\boldsymbol{f}_d)\, \mathrm{d}\mu_d,$$

with $\Delta_{\mathrm{ang}}^{-1}(I) = \{\boldsymbol{f}_d \in \mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}) \mid \Delta_{\mathrm{ang}}(Z(\boldsymbol{f}_d)) \in I\}$. This is a consequence of the coarea formula.

Example 4.14. Let $\boldsymbol{f}_d = (f_{1,d}, \dots, f_{n,d})$ be a random system of Laurent polynomials with $\operatorname{supp}(f_{i,d}) \subset dQ_i$ whose coefficients $\{f_{i,d,\boldsymbol{a}}\}_{\boldsymbol{a}\in\mathcal{A}_{i,d}}$ are independent complex Gaussian random variables with mean 0 and variance 1. This is a probability distribution on $\mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$ whose density function is defined, for $\boldsymbol{f}_d \in \mathbb{C}^{\boldsymbol{\mathcal{A}}_d}$, as

$$G_d(\boldsymbol{f}_d) = \prod_{i=1}^{n} \prod_{\boldsymbol{a}\in\mathcal{A}_{i,d}} \frac{1}{\pi} \mathrm{e}^{-|f_{i,d,\boldsymbol{a}}|^2} = \frac{1}{\pi^{\#\boldsymbol{\mathcal{A}}_d}} \mathrm{e}^{-\|\boldsymbol{f}_d\|_2^2} .$$

The random cycle $Z(\boldsymbol{f}_d)$ might be described by a probability distribution on $\mathbb{P}(\mathbb{C}^{\boldsymbol{\mathcal{A}}})$. The corresponding density function is the constant function $g_d = 1$. This can be seen by computing the integral along the fibers (4.13), or simply by observing that G_d is a function of the radius $\|\boldsymbol{f}_d\|_2$.

Theorem 1.8 implies then that the sequence of roots of $\boldsymbol{f}_d$ converge weakly to the equidistribution on $(S^1)^n$ when $d \to \infty$. In this way, we recover a result of Bloom and Shiffman [BS07, Example 3.5].

References

[AM96] F. Amoroso and M. Mignotte, *On the distribution of the roots of polynomials*, Ann. Inst. Fourier (Grenoble) **46** (1996), 1275–1291.

[Ber75] D. N. Bernstein, *The number of roots of a system of equations*, Funkcional. Anal. i Prilozen. **9** (1975), 1–4, English translation: Functional Anal. Appl. **9** (1975), 183–185.

[Bil97] Y. Bilu, *Limit distribution of small points on algebraic tori*, Duke Math. J. **89** (1997), 465–476.

[Bou70] N. Bourbaki, *Éléments de mathématique. Algèbre. Chapitres 1 à 3*, Hermann, 1970.

[BP07] C. Beltrán and L. M. Pardo, *Estimates on the distribution of the condition number of singular matrices*, Found. Comput. Math. **7** (2007), 87–134.

[BS07] T. Bloom and B. Shiffman, *Zeros of random polynomials on* $\mathbb{C}^m$, Math. Res. Lett. **14** (2007), 469–479.

[CLO05] D. A. Cox, J. Little, and D. O'Shea, *Using algebraic geometry*, second ed., Grad. Texts in Math., vol. 185, Springer, 2005.

[DS12] C. D'Andrea and M. Sombra, *An alternative definition of sparse resultants*, Preprint, 2012.

[ET50] P. Erdös and P. Turán, *On the distribution of roots of polynomials*, Ann. of Math. (2) **51** (1950), 105–119.

[FRL06] C. Favre and J. Rivera-Letelier, *Équidistribution quantitative des points de petite hauteur sur la droite projective*, Math. Ann. **335** (2006), 311–361, *Corrigendum* in Math. Ann. **339** (2007), 799–801.

[Gan54] T. Ganelius, *Sequences of analytic functions and their zeros*, Ark. Mat. **3** (1954), 1–50.

[GKZ94] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Birkhäuser, 1994.

[HN08] C. P. Hughes and A. Nikeghbali, *The zeros of random polynomials cluster uniformly near the unit circle*, Compos. Math. **144** (2008), 734–746.

[Kho91] A. G. Khovanskii, *Fewnomials*, Transl. Math. Monogr., vol. 88, Amer. Math. Soc., 1991.

[KN69] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Interscience Tracts in Pure and Applied Mathematics, No. 15, John Wiley & Sons, 1969.

[Min03] M. Minimair, *Sparse resultant under vanishing coefficients*, J. Algebraic Combin. **18** (2003), 53–73.

[Pet05] C. Petsche, *A quantitative version of Bilu's equidistribution theorem*, Int. J. Number Theory **1** (2005), 281–291.

[PS93] P. Pedersen and B. Sturmfels, *Product formulas for resultants and Chow forms*, Math. Z. **214** (1993), 377–396.

[Som04] M. Sombra, *The height of the mixed sparse resultant*, Amer. J. Math. **126** (2004), 1253–1260.

[SZ04] B. Shiffman and S. Zelditch, *Random polynomials with prescribed Newton polytope*, J. Amer. Math. Soc. **17** (2004), 49–108.

D'Andrea: Universitat de Barcelona, Departament d'Àlgebra i Geometria. Gran Via 585, 08007 Barcelona, Spain
E-mail address: `cdandrea@ub.edu`
URL: `http://atlas.mat.ub.es/personals/dandrea/`

Galligo: Université de Nice-Sophia Antipolis, Laboratoire de Mathématiques. Parc Valrose, 06108 Nice Cedex 02, France
E-mail address: `galligo@unice.fr`
URL: `http://math.unice.fr/~galligo/`

Sombra: ICREA and Universitat de Barcelona, Departament d'Àlgebra i Geometria. Gran Via 585, 08007 Barcelona, Spain
E-mail address: `sombra@ub.edu`
URL: `http://atlas.mat.ub.es/personals/sombra/`